

# Activation of Protein Kinase C $\beta$ II by the Stereo-specific Phosphatidylserine Receptor Is Required for Phagocytosis of Apoptotic Thymocytes by Resident Murine Tissue Macrophages\*

Received for publication, March 27, 2002, and in revised form, June 25, 2002  
Published, JBC Papers in Press, July 11, 2002, DOI 10.1074/jbc.M202967200

Jill C. Todt<sup>‡</sup>, Bin Hu<sup>‡</sup>, Antonello Punturieri<sup>‡§</sup>, Joanne Sonstein<sup>‡</sup>, Timothy Polak<sup>‡</sup>,  
and Jeffrey L. Curtis<sup>‡§¶\*\*</sup>

From the <sup>‡</sup>Division of Pulmonary and Critical Care Medicine, Department of Internal Medicine, the <sup>¶</sup>Comprehensive Cancer Center, and the <sup>§</sup>Graduate Program in Immunology, University of Michigan Health Care System and the <sup>§</sup>Pulmonary and Critical Care Medicine Section, Medical Service, Department of Veterans Affairs Care System, Ann Arbor, Michigan 48105-2303

We showed previously that protein kinase C (PKC) is required for phagocytosis of apoptotic leukocytes by murine alveolar (AM $\phi$ ) and peritoneal macrophages (PM $\phi$ ) and that such phagocytosis is markedly lower in AM $\phi$  compared with PM $\phi$ . In this study, we examined the roles of individual PKC isoforms in phagocytosis of apoptotic thymocytes by these two M $\phi$  populations. By immunoblotting, AM $\phi$  expressed equivalent PKC  $\eta$  but lower amounts of other isoforms ( $\alpha$ ,  $\beta$ I,  $\beta$ II,  $\delta$ ,  $\epsilon$ ,  $\mu$ , and  $\zeta$ ), with the greatest difference in  $\beta$ II expression. A requirement for PKC  $\beta$ II for phagocytosis was demonstrated collectively by phorbol 12-myristate 13-acetate-induced depletion of PKC  $\beta$ II, by dose-response to PKC inhibitor Ro-32-0432, and by use of PKC  $\beta$ II myristoylated peptide as a blocker. Exposure of PM $\phi$  to phosphatidylserine (PS) liposomes specifically induced translocation of PKC  $\beta$ II and other isoforms to membranes and cytoskeleton. Both AM $\phi$  and PM $\phi$  expressed functional PS receptor, blockade of which inhibited PKC  $\beta$ II translocation. Our results indicate that murine tissue M $\phi$  require PKC  $\beta$ II for phagocytosis of apoptotic cells, which differs from the PKC isoform requirement previously described in M $\phi$  phagocytosis of other particles, and imply that a crucial action of the PS receptor in this process is PKC  $\beta$ II activation.

Phagocytosis, the uptake of large particles ( $>0.5\ \mu\text{m}$ ) via actin-dependent mechanisms (1), is the obligatory means of clearing apoptotic cells during development and in resolving inflammation (2). Only macrophages can efficiently clear the large numbers of apoptotic leukocytes produced during waning immune responses (3–6). Indeed, the efficiency of this process is evidenced by the fact that apoptotic cells are rarely observed *in vivo* (7); one exception is in the lungs of mice, where apoptotic lymphocytes are easily demonstrable both in health and

inflammation (8). This defect in clearance is consistent with the finding that the principal resident lung phagocytes, alveolar macrophages (AM $\phi$ ),<sup>1</sup> exhibit markedly lower capacity for phagocytosis of apoptotic leukocytes, either compared with inflammatory lung M $\phi$  (in rabbits) (9) or with resident peritoneal M $\phi$  (PM $\phi$ ) (in mice) (10). In the latter system, no disparity between AM $\phi$  and PM $\phi$  was detected using three other particle types (10, 11). Murine AM $\phi$  also exhibited a relative deficit in phagocytosis of apoptotic cells *in vivo* (10). We have recently found that human AM $\phi$  also show much lower phagocytosis of apoptotic cells than of other particles *in vitro*.<sup>2</sup> Contrasting the properties of these two types of resident tissue M $\phi$  could aid in defining the molecular basis of apoptotic cell recognition, which is poorly understood.

Recognition of apoptotic cells is initiated through at least two pathways. Using a 70-kDa glycosylated type II transmembrane protein called PS-R' (12), M $\phi$  and other cell types recognize externalized phosphatidylserine (PS), which translocates from the inner to the outer leaflet of the cell membrane early in apoptosis (13–17). Recognition of externalized PS has been suggested to be both necessary and sufficient to generate a signal for ingestion (13, 18). More recently, the M $\phi$ -specific receptor tyrosine kinase Mer has been identified as critical for the phagocytosis of apoptotic cells by murine M $\phi$  (19). How signaling from these two receptors leads to apoptotic cell phagocytosis is undefined. A host of other M $\phi$  cell surface receptors (reviewed in Ref. 20) have also been implicated in clearance of apoptotic cells, but they appear to be involved principally in adhesion rather than in recognition of cell death (21). Moreover, although we (10) and others (22) have identified a number of differences in expression of adhesion molecules between murine AM $\phi$  and PM $\phi$ , blocking experiments using monoclonal antibodies (mAbs) or the arginine-glycine-aspartic acid-serine (RGDS) tetrapeptide have not supported any identified adhesion receptor, including several integrins, as being responsible for the functional difference in phagocytosis (10).

An alternative explanation for disparity between M $\phi$  types in apoptotic cell phagocytosis would be differences in postreceptor signal transduction. A logical candidate for such a difference is

\* This work was supported by United States Public Health Service Grants RO1 HL56309 and RO-1 HL6157, Merit Review funding, and a Research Enhancement Award Program grant from the Department of Veterans Affairs. Portions of this work have been presented previously at the Autumn Immunology Conference, Chicago, IL, November 17, 2001, and at the American Thoracic Society Meeting in Atlanta, GA, May 22, 2002. The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked "advertisement" in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

\*\* To whom correspondence should be addressed: Pulmonary and Critical Care Medicine Section (506/111G), Dept. of Veterans Affairs Medical Center, 2215 Fuller Rd., Ann Arbor, MI 48105-2303. Tel.: 734-761-7980; Fax: 734-761-7843; E-mail: jlc Curtis@umich.edu.

<sup>1</sup> The abbreviations used are: AM $\phi$ , alveolar macrophage(s); aPKC, atypical PKC isoform; cPKC, conventional PKC isoform; DAG, diacylglycerol; mAb, monoclonal antibody; M $\phi$ , macrophage(s); nPKC, novel PKC isoform; PI, phosphatidylinositol; PKC, protein kinase C; PM $\phi$ , peritoneal macrophages; PMA, phorbol 12-myristate 13-acetate; PS, phosphatidylserine; PBS, phosphate-buffered saline; RACK, receptor for activated C kinase.

<sup>2</sup> J. C. Todt, B. Hu, A. Punturieri, J. Sonstein, T. Polak, and J. L. Curtis, unpublished observation.

protein kinase C (PKC), because we and others have shown that it is required for apoptotic cell clearance (11, 23). PKC comprises a family of related serine/threonine kinases divided into three groups on the basis of structure and cofactor requirements (24). Activation of PKC requires phosphorylation on serines/threonines, displacement of its autoinhibitory pseudosubstrate domain, and translocation to specific cytoskeletal and intracellular membrane sites of action (25). Activation of the conventional group (cPKC) ( $\alpha$ ,  $\beta$ I,  $\beta$ II, and  $\gamma$ ) is calcium- and diacylglycerol (DAG)-dependent. Activation of the novel group (nPKC) ( $\epsilon$ ,  $\delta$ ,  $\eta$ , and  $\theta$ ) also depends on binding of DAG, but it is calcium-independent. The atypical group (aPKC) ( $\iota$ / $\lambda$  and  $\zeta$ ) cannot be activated by calcium or DAG. All PKC family members bind PS on the cytosolic leaflet of the cell membrane, but aPKCs require additional incompletely defined lipid activators (24). Another isoform, PKC  $\mu$  (often called PKD in the mouse), does not fit into any of the major groups. PKC  $\mu$  contains two unique hydrophobic domains in its amino terminus and is phospholipid-dependent but calcium-insensitive (26). Individual cell types usually express several PKC isoforms, each of which appears to mediate unique functions (27). Even the 50-amino acid difference in the alternatively spliced forms of PKC  $\beta$  ( $\beta$ I and  $\beta$ II) appears to be responsible for the unique role of each PKC  $\beta$  isozyme (28, 29). Thus, differences between murine AM $\phi$  and PM $\phi$  in PKC isoform expression or function could explain the functional difference between these two cell types in apoptotic cell phagocytosis.

We recently showed (11) that phagocytosis of apoptotic thymocytes by murine AM $\phi$  and PM $\phi$  was reduced by the nonspecific PKC inhibitor staurosporine and by Gö 6976 but only incompletely by calphostin C. Gö 6976 has been reported to act as a partially selective inhibitor of the cPKC  $\alpha$  and  $\beta$ I isoforms (30), whereas calphostin C has greater activity against nPKCs than cPKCs (31). However, current data on the specificity of these inhibitors are too inconclusive (32) to allow us to predict with certainty which isoforms are involved. Therefore, in this study, we used six approaches to further define the PKC isoform(s) involved in M $\phi$  phagocytosis of apoptotic thymocytes. First, because the pattern of PKC isoforms in primary murine tissue M $\phi$  has not been described, we analyzed PKC isoform expression using isotype-specific antibodies and Western blotting. Second, we tested the effect of overnight exposure of M $\phi$  to phorbol 12-myristate 13-acetate (PMA), which depletes cPKC and nPKC isoforms by interacting with their DAG-binding sites, on AM $\phi$  and PM $\phi$  phagocytosis of apoptotic thymocytes. Third, we employed the isoform-selective inhibitors, rottlerin and Ro-32-0432. Fourth, we tested the effect of myristoylated blocking peptides against PKC  $\beta$ I, PKC  $\beta$ II, and PKC  $\alpha$  on phagocytosis of apoptotic thymocytes by AM $\phi$  and PM $\phi$ . Fifth, we examined the effect of PS liposomes, as models for apoptotic thymocytes, on translocation of PKC isoforms to cell membranes and cytoskeleton; apoptotic thymocytes could not be used, because they themselves express multiple PKC isoforms. Finally, we studied M $\phi$  expression of PS-R' and the relationship between PS-R' stimulation and PKC translocation. Collectively, our results indicate a requirement for PKC  $\beta$ II in M $\phi$  phagocytosis of apoptotic thymocytes, show that an antibody against PS-R' blocks translocation of PKC  $\beta$ II in response to PS liposomes, and suggest that relative deficiency of PKC  $\beta$ II and possibly other PKC isoforms may partially explain the functional difference in apoptotic cell clearance by AM $\phi$ .

#### EXPERIMENTAL PROCEDURES

**Reagents**—Rottlerin and Ro-32-0432 were purchased from Calbiochem. PMA, PBS, RPMI 1640, fetal bovine serum, HEPES, pyruvate, 1 kb Plus brand up markers and penicillin/streptomycin were obtained from Invitrogen. Dimethyl sulfoxide, dexamethasone, 2-mercaptoethanol, sodium deoxycholate, glycerol, NaCl, Tris-HCl, Triton X-100,

Tween 20, 1 kb Plus brand up markers, L- $\alpha$ -phosphatidylinositol (PI), L- $\alpha$ -PS, and phosphatase inhibitor mixture II were purchased from Sigma. Antibodies and blocking peptides for PKC isoforms and horseradish peroxidase-conjugated anti-rabbit IgG were purchased from Santa Cruz Biotechnology, Inc. (Santa Cruz, CA). Complete mini-protease inhibitor tablets and the lactate dehydrogenase cytotoxicity detection kit were purchased from Roche Molecular Biochemicals. SDS, 0.2- $\mu$ m polyvinylidene difluoride membrane, nonfat dry milk blocker, and 10% Ready Acrylamide gels were obtained from Bio-Rad. Supersignal West Femto Maximum Sensitivity substrate was obtained from Pierce. Eastman Kodak Co. X-Omat AR film and eight-well Lab-Tek slides were obtained from Fisher. mAb 217 (anti-murine PS-R; rat IgM) was generously provided by Dr. Valerie Fadok (National Jewish Medical Center, Denver, CO) as a culture supernatant; in selected experiments, a commercial preparation of this mAb (Cascade BioScience, Winchester, MA) was used. Myristoylated PKC peptides were synthesized by BIOSOURCE Quality Controlled Biochemicals (Hopkinton, MA) at 90% purity, as confirmed by high pressure liquid chromatography and mass spectroscopy performed by the manufacturer.

**Mice**—All experiments were performed using pathogen-free C57BL/6 female mice purchased from Charles River Laboratories Inc. (Wilmington, MA) at 7–8 weeks of age and used at 8–14 weeks of age. Mice were housed in the Animal Care Facility at the Ann Arbor VA Medical Center, which is fully accredited by the American Association for Accreditation of Laboratory Animal Care. This study complied with the National Institutes of Health "Guide for the Care and Use of Laboratory Animals" (Department of Health, Education, and Welfare Publication No. 80-32) and followed a protocol approved by the Animal Care Subcommittee of the local institutional review board.

**Isolation and Culture of M $\phi$** —Mice were euthanized by asphyxia in a high CO<sub>2</sub> environment, which we have previously shown does not impair the capacity of AM $\phi$  to ingest apoptotic thymocytes (10). Resident AM $\phi$  and PM $\phi$  were harvested and cultured as previously described (10). M $\phi$  were isolated by adherence onto tissue culture plates (for protein isolation) or eight-well Lab-Tek slides (for phagocytosis assay) for 2–4 h at 37 °C in 5% CO<sub>2</sub> in complete medium (RPMI 1640 containing 10% heat-inactivated fetal bovine serum, 25 mM HEPES, 2 mM L-glutamine, 1 mM pyruvate, 100 units/ml penicillin/streptomycin, 55  $\mu$ M 2-mercaptoethanol). Nonadherent cells were removed by gentle washing. In experiments in which PKC localization was analyzed by Western blotting, complete medium was replaced with serum-free medium for 1 h.

**Preparation of Apoptotic Thymocytes**—Thymuses were harvested from normal mice and minced to yield a single cell suspension. To induce apoptosis, thymocytes were resuspended in complete medium to a concentration of  $1 \times 10^6$  cells/ml and incubated for 6 h in complete medium containing 1  $\mu$ M dexamethasone. This treatment yields a population with a low percentage (mean 13.4%) contamination of late apoptotic or necrotic cells (10, 11).

**Western Analysis of PKC Isozymes**—Resident AM $\phi$  and PM $\phi$  from normal mice were isolated and plated at  $1.0 \times 10^6$  cells/ml on 100  $\times$  15-mm plates in medium containing 10% serum and incubated for 2 h at 37 °C and 5% CO<sub>2</sub>. Next, M $\phi$  were washed and solubilized in ice-cold lysis buffer consisting of 1.0% Triton X-100, 20 mM Tris-HCl, pH 8.0, 150 mM NaCl, 0.1% SDS, 0.5% sodium deoxycholate, and 10% glycerol with protease inhibitors (complete minitab) and phosphatase inhibitor mixture II (1:100) for 30 min on ice. After sonicating for 3 s and centrifuging at  $13,800 \times g$  for 3 min, 7  $\mu$ g of protein/sample was run on a 10% acrylamide gel under reducing conditions and transferred to a polyvinylidene difluoride membrane using 25 mM Tris, 192 mM glycine, 20% methanol. Blots were blocked with 5% nonfat dry milk in PBS (blocker), incubated with the appropriate anti-PKC isoform antibody (1:1000 dilution in blocker), and washed five times for 5 min each with PBS containing 0.1% Tween 20. Blots were incubated with horseradish peroxidase-conjugated goat anti-rabbit IgG (diluted 1:10,000 in PBS containing 5% nonfat dry milk), and the chemiluminescence signal was developed by adding a peroxidase/luminol-based substrate (Supersignal Femto reagent; Pierce). The identity of PKC isoforms on Western blots was verified using isoform-specific blocking peptides. No bands were seen when blots were stained using horseradish peroxidase-conjugated goat anti-rabbit IgG alone.

To analyze the subcellular distribution of PKC isozymes, M $\phi$  were isolated and plated at  $1.0 \times 10^6$  cells/ml on 30  $\times$  15-mm plates in medium containing 10% serum and incubated for 2 h at 37 °C and 5% CO<sub>2</sub>. The medium was then removed and replaced with serum-free medium for 1 h followed by washing and scraping in cold PBS. Next, M $\phi$  were sonicated for 10 s in 50 mM Tris, pH 8.0, 9 mM EDTA and then centrifuged at  $100,000 \times g$  for 45 min. The supernatant was collected



and defined as the cytosolic fraction. The pellet, defined as the membrane and cytoskeleton fractions, was solubilized in ice-cold lysis buffer (defined above) for 30 min on ice. This solubilized pellet fraction was sonicated for 3 s and then centrifuged at  $13,800 \times g$  for 3 min. Equal amounts of protein (3–7  $\mu$ g) were run on a 10% SDS-PAGE gel under reducing conditions, transferred to the polyvinylidene difluoride membrane, and stained as described above.

**Preparation of Liposomes**—To produce liposomes, PS or the negatively charged control lipid PI was dried under  $N_2$  and resuspended in serum-free medium by vortexing. Liposome size was determined by Coulter counter analysis to be in a similar range as apoptotic thymocytes, 2–3.2  $\mu$ m. In these experiments, M $\phi$  were incubated for 1 h in serum-free medium, liposomes in serum-free medium were added to M $\phi$  monolayers in a final PS or PI concentration of 0.11 mM, and M $\phi$  and liposomes were co-incubated for the indicated time at 37 °C and 5%  $CO_2$ . Cells were washed and scraped in cold PBS; sonicated for 10 s in 50 mM Tris, pH 8.0, 9 mM EDTA; and centrifuged at  $100,000 \times g$  for 45 min.


**RNA Preparation and Reverse Transcriptase-PCR**—Total RNA was isolated from adherent AM $\phi$  and PM $\phi$  using TRIzol (Invitrogen). Reverse transcriptase-PCRs were performed using kits from Invitrogen. The primer sets used were the following: for mouse PS-R' (GenBank<sup>TM</sup> accession number AF304118), forward CTC ACG ATG AAC CAC AAG AGC and reverse GGA CCA GCC CTC TTG TGC ATT; for mouse glyceraldehyde-3-phosphate dehydrogenase (GenBank<sup>TM</sup> accession number M32599), forward GGT CGG TGT GAA CGG ATT TGG and reverse ATG AGG TCC ACC ACC CTG TTG. The expected PCR product sizes are 245 bp for PS-R' and 968 bp for glyceraldehyde-3-phosphate dehydrogenase. PCR products were analyzed on a 2% agarose gel and stained with ethidium bromide. The identity of the target products was confirmed by sequencing.

**Flow Cytometric Analysis**—Apoptosis was measured by simultaneous annexin V and propidium iodide staining (apoptosis detection kit; R & D Systems, Minneapolis, MN) according to the manufacturer's protocol. Cells were analyzed without fixation by flow cytometry within 1 h of staining. Staining of surface receptors and flow cytometry were performed as previously described in detail (33) using a FACScan cytometer (BD PharMingen) running Cell Quest software on a PowerPC microcomputer (Apple, Cupertino, CA) for data collection and analysis. A minimum of 10,000 cells were analyzed.

**Phagocytosis Assay**—Phagocytosis of apoptotic thymocytes *in vitro* was assayed by co-incubation of  $0.5\text{--}2.0 \times 10^5$  adherent M $\phi$  with  $2.0 \times 10^6$  apoptotic thymocytes for 90 min at 37 °C in 5%  $CO_2$  as previously described (10). Results are expressed as the percentage of M $\phi$  containing at least one ingested thymocyte (percentage of phagocytosis) and as the phagocytic index, which was generated by multiplying the percentage of phagocytosis by the mean number of ingested cells per M $\phi$ . Cell-permeable PKC inhibitors were added either 30 min (rottlerin) or 18 h (Ro-32-0432 and PMA) before the addition of apoptotic thymocytes. Myristoylated PKC peptides from the carboxyl terminus of the V5 region of PKC  $\alpha$  (myr-PQFVHPILQSAV-amide), PKC  $\beta$ I (myr-DQNE-FAGFSYTNPEFVINV-amide), or PKC  $\beta$ II (myr-SFVNSEFLKPEVKS-amide) were added to a final concentration of 100  $\mu$ M 30 min before the addition of apoptotic thymocytes. The myristate moieties coupled to the amino terminus of these peptides allow membrane permeability, permitting their use in primary cells (34). All inhibitors were nontoxic at the times and concentrations utilized, as determined by the lactate dehydrogenase cytotoxicity detection kit.<sup>2</sup>

**Adhesion Assay**—Adherence of apoptotic thymocytes to M $\phi$  *in vitro* was assayed in the same fashion as phagocytosis, except that  $2 \times 10^7$  apoptotic thymocytes suspended in 400  $\mu$ l of complete medium were added to each well, yielding a 100:1 ratio of thymocytes to M $\phi$ . The slides were incubated for various times at 37 °C and then washed in a standardized fashion by dipping individual slides five times in each of two Wheaton jars filled with ice-cold PBS. Slides were then stained using hematoxylin-eosin Y (Richard-Allan; Kalamazoo, MI) and cover-slipped. Adhesion was evaluated by counting 200–300 M $\phi$  per well at  $\times 1000$  magnification under oil immersion and scoring for bound thymocytes. Results are expressed as the percentage of M $\phi$  binding at least one thymocyte (percentage of adhesive M $\phi$ ) and as the adhesion index, which was generated by multiplying the percentage of adherence-positive M $\phi$  by the mean number of adherent thymocytes per M $\phi$ . In blocking experiments, mAbs were used at concentrations previously determined to be saturating by flow cytometry, and culture supernatant containing mAb 217 was concentrated 10-fold using Centrprep tubes (Amicon, Bedford, MA).

**Statistical Analysis**—Data are expressed as mean  $\pm$  S.E. Statistical calculations were performed using Statview on a Macintosh Power PC




**FIG. 1. Resident murine AM $\phi$  and PM $\phi$  differ in expression of multiple PKC isoforms.** Whole cell lysates of AM $\phi$  and PM $\phi$  of normal C57BL/6 mice cultured as adherent monolayers for 2 h were analyzed for expression of PKC isoforms by Western blotting using isoform-specific antibodies.

G3 computer. Continuous ratio scale data were evaluated by unpaired Student's *t* test (for two samples) or analysis of variance (for multiple comparisons) with *post hoc* analysis by the Tukey-Kramer test or by the two-tailed Dunnett test, which specifically compares treatment groups with a control group (35). Use of these parametric statistics was deemed appropriate, since phagocytosis of apoptotic thymocytes by PM $\phi$  has been shown to follow a Gaussian distribution (36). Significant differences were defined as  $p < 0.05$ .

## RESULTS

**AM $\phi$  Express Lower Amounts of Most PKC Isoforms than Do PM $\phi$** —Western blot analysis using PKC isoform-specific antibodies demonstrated markedly lower expression of PKC  $\alpha$ ,  $\beta$ I,  $\beta$ II,  $\delta$ ,  $\epsilon$ ,  $\mu$ , and  $\xi$  in resident murine AM $\phi$  in comparison with resident murine PM $\phi$  (Fig. 1). By contrast, AM $\phi$  had slightly higher expression of PKC  $\eta$ . Staining did not detect expression of PKC  $\gamma$  or PKC  $\lambda$  in either murine AM $\phi$  or PM $\phi$ ,<sup>2</sup> in agreement with previous analyses of human tissue M $\phi$  (24, 37), and we did not test expression of the lymphocyte-specific isoform PKC  $\theta$ . Thus, the lower expression of several PKC isoforms by AM $\phi$  is a potential explanation for the previously observed decreased phagocytosis of apoptotic cells by this cell type.


**Effect of Overnight PMA Treatment on M $\phi$  Phagocytosis of Apoptotic Thymocytes**—Chronic (18-h) PMA treatment depletes cPKCs and nPKCs but not aPKCs, with the degree to which individual isoforms are affected depending on the cell type and the PMA concentration (27, 37–39). In preliminary experiments, the concentration and time of PMA addition were optimized to achieve maximal inhibition of apoptotic cell phagocytosis. We determined that the conditions used were nontoxic, as indicated by assay of lactate dehydrogenase release.<sup>2</sup> Western blot analysis on AM $\phi$  and PM $\phi$  after overnight treatment using 8.1  $\mu$ M PMA confirmed that PKC  $\alpha$ ,  $\beta$ I,  $\beta$ II, and  $\delta$  were significantly depleted ( $p < 0.05$ , unpaired *t* test) by PMA treatment in either of the two types of M $\phi$  (Fig. 2). As anticipated, PKC  $\epsilon$ ,  $\eta$ ,  $\mu$ , and  $\zeta$  were not depleted by overnight PMA




**FIG. 2. Overnight PMA treatment decreases expression of cPKC and nPKC isoforms by AM $\phi$  and PM $\phi$ .** Resident AM $\phi$  and PM $\phi$  were incubated overnight at 37 °C and 5% CO<sub>2</sub> with either PMA (8.1  $\mu$ M final concentration) or the appropriate amount of ethanol as the control and then analyzed for expression of PKC isoforms by Western blotting using isoform-specific antibodies.

treatment in either M $\phi$  type. In the functional assay, the same overnight PMA treatment significantly decreased phagocytosis by AM $\phi$  and PM $\phi$  ( $p < 0.05$ , unpaired Student  $t$  test) (Fig. 3). Control experiments showed that overnight PMA treatment did not influence adhesion of thymocytes to PM $\phi$  (percentage of adhesive M $\phi$ , 86.1  $\pm$  1.7% (control) versus 82.0  $\pm$  2.0% (PMA-treated),  $p = 0.16$ ; adhesion index, 2.6  $\pm$  0.2 (control) versus 2.7  $\pm$  0.1 (PMA-treated); mean  $\pm$  S.E.,  $n = 4$ ,  $p = 0.47$ , both comparisons made by unpaired  $t$  test) and actually slightly increased adhesion to AM $\phi$  (adhesive M $\phi$ , 60.5  $\pm$  1.5% (control) versus 71.5  $\pm$  2.2% (PMA-treated); mean S.E.,  $n = 4$ ;  $p < 0.001$ ; adhesion index, 1.2  $\pm$  0.1 (control) versus 1.7  $\pm$  0.0 (PMA-treated);  $p < 0.001$ , unpaired  $t$  test). Collectively, these data indicate that the nPKC isoforms  $\epsilon$  and  $\eta$ , the aPKC isoform  $\zeta$ , and PKC  $\mu$ /PKD are not essential for M $\phi$  phagocytosis of apoptotic cells but leave open the question of which cPKC or other nPKC isoforms are required.

**Rottlerin and Ro-32-0432 Decrease Phagocytosis of Apoptotic Thymocytes at Concentrations at Which They Inhibit cPKCs**—To further define which PKC isoforms are essential for phagocytosis, we used two cell-permeable PKC inhibitors, rottlerin and Ro-32-0432, which show a degree of isoform specificity. Rottlerin specifically inhibits PKC  $\delta$  at relatively low concentration (IC<sub>50</sub> = 3–6  $\mu$ M) and other PKC isoforms only at higher concentrations (IC<sub>50</sub> for PKC  $\alpha$  and  $\beta$  = 30–42  $\mu$ M; IC<sub>50</sub> for PKC  $\epsilon$ ,  $\eta$ , and  $\zeta$  is 80–100  $\mu$ M) (40). We found that rottlerin did not inhibit AM $\phi$  or PM $\phi$  phagocytosis of apoptotic thymocytes at 10  $\mu$ M (a concentration at which PKC  $\delta$  should be inhibited (40)), whereas there was significantly decreased phagocytosis at 40  $\mu$ M (a concentration at which cPKCs should be inhibited (40)) (Fig. 4). Ro-32-0432 had no effect on phagocytosis of apoptotic thymocytes at 9 nM (a concentration at which PKC  $\alpha$  should be inhibited (41)) but did inhibit phagocytosis of apoptotic thymocyte at 28 nM (a concentration at which PKC  $\beta$  but not PKC  $\epsilon$  should be inhibited (41)) (Fig. 5). These data argue against a requirement for PKC  $\delta$  or PKC  $\epsilon$  for phagocytosis of apoptotic cells and provide additional support for the possibility that a cPKC, especially PKC  $\beta$ I or  $\beta$ II, is required. However, rottlerin can inhibit casein kinase II and cAMP-dependent protein kinase with IC<sub>50</sub> values of 30 and 78  $\mu$ M, respectively, and can also inhibit calmodulin kinase III with IC<sub>50</sub> of 5.3  $\mu$ M (32). Moreover, interpretation of these experiments is complicated by the uncertainty about the



**FIG. 3. Depletion of cPKCs and nPKCs by overnight PMA treatment reduces M $\phi$  phagocytosis of apoptotic thymocytes.** Phagocytosis of apoptotic thymocytes by resident AM $\phi$  and PM $\phi$  incubated overnight in medium containing a final concentration of 8.1  $\mu$ M PMA (black bars) or the appropriate amount of ethanol as a control (gray bars) was determined by examining hematoxylin-eosin Y-stained slides under oil immersion. A, percentage of phagocytic M $\phi$ ; B, phagocytic index. Data are mean  $\pm$  S.E. of four replicates in two experiments. \*,  $p < 0.05$ , unpaired Student's  $t$  test.



**FIG. 4. Rottlerin inhibits M $\phi$  phagocytosis of apoptotic thymocytes in a dose-dependent fashion.** Phagocytosis of apoptotic thymocytes by resident AM $\phi$  and PM $\phi$  was determined after a 30-min incubation under the following conditions: control medium (light gray bars), 10  $\mu$ M rottlerin (black bars), 40  $\mu$ M rottlerin (white bars), and 100  $\mu$ M rottlerin (dark gray bars). A, percentage of phagocytic M $\phi$ ; B, phagocytic index. Data are mean  $\pm$  S.E. of three replicates and are representative of results of two experiments. \*,  $p < 0.05$ , analysis of variance with *post hoc* Dunnett's testing.

degree to which the reported IC<sub>50</sub> values for these inhibitors can be extrapolated from the *in vitro* assays with which they were developed to intact cells. Therefore, we looked for an




FIG. 5. **Ro-32-0432 inhibits Mφ phagocytosis of apoptotic thymocytes in a dose-dependent fashion.** Phagocytosis of apoptotic thymocytes by resident AMφ and PMφ was determined after 30 min of incubation under the following conditions: control medium (gray bars), 9  $\mu$ M Ro-32-0432 (black bars), 28  $\mu$ M Ro-32-0432 (white bars). A, percentage of phagocytic Mφ; B, phagocytic index. Data are mean  $\pm$  S.E. of four replicates in two experiments. \*,  $p < 0.05$ , analysis of variance with *post hoc* Dunnett's testing.

additional method to verify a requirement of PKC  $\beta$  during phagocytosis.

**Myristoylated PKC  $\beta$ II Blocking Peptide Inhibits Phagocytosis of Apoptotic Thymocytes by AMφ and PMφ**—Because all data thus far implicated a cPKC as a necessary signaling component during phagocytosis of apoptotic thymocytes, we tested the effect of cell-permeable myristoylated peptides derived from the carboxyl terminus of the V5 region of PKC  $\alpha$ ,  $\beta$ I, or  $\beta$ II, on phagocytosis of apoptotic thymocytes. Peptides from this region were selected based on the following rationale. First, this region (specifically the last 13 amino acids) has been suggested to contribute to phosphatidylglycerol-induced activation of PKC  $\beta$ II (29) and, via interaction with the C2 region, to calcium- and PS-mediated activation of PKC  $\beta$ II (42). Second, the V5 region of PKC  $\alpha$ , specifically the QSAV sequence, has been shown to interact with PICK1, a PKC-binding protein that appears to target PKC  $\alpha$  to appropriate intracellular sites for transduction of isozyme-specific signals (43). Third, the PKC  $\beta$ II peptide used in this study has previously been shown to inhibit the binding of a maltose-binding protein-PKC  $\beta$ II fusion protein to RACK1 (34), confirming its functional activity. Fourth, the V5 region is the only site at which PKC  $\beta$ I and PKC  $\beta$ II amino acid sequences differ. Finally, these  $\beta$ I and  $\beta$ II peptides comprise part of the antigenic sequences used for development of the isozyme-specific antibodies used in this study.

We found that the myristoylated PKC  $\beta$ II peptide significantly decreased phagocytosis of apoptotic thymocytes by both AMφ and PMφ, whereas the myristoylated PKC  $\alpha$  and PKC  $\beta$ I peptides had no effect (Fig. 6). These results provide direct evidence that PKC  $\beta$ II is required for Mφ phagocytosis of apoptotic thymocytes.

**PS Liposomes Stimulate Translocation of PKC Isoforms  $\beta$ I,  $\beta$ II,  $\delta$ ,  $\epsilon$ ,  $\mu$ , and  $\zeta$  to Membrane and Cytoskeleton Fractions**—Translocation from cytosol to membrane or cytoskeleton fractions upon activation is a major mechanism of PKC regulation (26, 44), leading us to investigate translocation of individual PKC isoforms during the phagocytic process. However, prelim-





FIG. 6. **Myristoylated PKC  $\beta$ II blocking peptide specifically inhibits Mφ phagocytosis of apoptotic thymocytes.** Phagocytosis of apoptotic thymocytes by resident AMφ and PMφ was determined after 30 min of incubation in control medium (light gray bars) or in medium containing a final concentration of 100  $\mu$ M in water of one of the following myristoylated blocking peptides: PKC  $\alpha$  peptide (black bars), PKC  $\beta$ I (white bars), or PKC  $\beta$ II peptide (dark gray bars). A, percentage of phagocytic Mφ; B, phagocytic index. Data are mean  $\pm$  S.E. of three replicates and are representative of two experiments.

inary experiments showed that the abundant expression of multiple PKC isoforms in viable murine thymocytes was retained upon induction of apoptosis, making use of apoptotic thymocytes for these studies unfeasible. Instead, we substituted PS liposomes, which have been used previously as models for apoptotic cells (45, 46). Liposome clearance by AMφ and PMφ has been shown to require phagocytosis and not to result from membrane fusion (47, 48), further supporting use of PS liposomes for these studies. Based on the practical consideration that our previous experiment showed that AMφ express much lower amounts of multiple PKC isoforms (Fig. 1), we performed these experiments using only PMφ. Time points were selected based on our previous study of the kinetics of phagocytosis (10).

We found that PS liposomes strongly stimulated translocation of PKC  $\beta$ I,  $\beta$ II,  $\delta$ ,  $\epsilon$ ,  $\mu$ , and  $\zeta$  from the cytosol to the membrane and cytoskeleton fraction of PMφ by 10 min, whereas control PI liposomes had no effect (Fig. 7). PS liposomes specifically stimulated slight translocation of PKC  $\alpha$ , and PI liposomes also slightly stimulated translocation of PKC  $\delta$  and  $\epsilon$ , whereas neither type of liposome had a significant effect on the localization of PKC  $\eta$ . Thus, several members of all three groups of PKC isoforms are specifically translocated in response to PS exposure.

**Resident Murine Mφ Express the Stereo-specific PS-R', Which Mediates PKC Translocation in Response to PS Liposomes**—These data raised the question of whether the stereo-specific PS-R' identified by Fadok and associates (12) could contribute to PKC activation during Mφ recognition of apoptotic cells. Expression of PS-R' was detected by multiple methods in both types of Mφ. Reverse transcriptase-PCR demonstrated equivalent PS-R' mRNA in AMφ and PMφ (Fig. 8A). Flow cytometry showed distinct surface expression of PS-R' by both types of Mφ (Fig. 8, B and C). Although specific staining and background staining of both Mφ types varied slightly between experiments,






**FIG. 7. PS liposomes induce translocation of multiple PKC isoforms into the membrane and cytoskeletal fractions.** Resident murine PM $\phi$  were incubated for various times with liposomes consisting of L- $\alpha$ -PS (left-hand column) or control L- $\alpha$ -PI (right-hand column), and then expression of PKC isoforms in the cytosolic fraction and in the membrane and cytoskeletal fraction was analyzed by Western blotting. C, control (no liposomes).

mean channel fluorescence of PM $\phi$  was generally somewhat higher than that of AM $\phi$  (e.g. in the experiment shown in Fig. 8, mean channel fluorescence was 484.7 for PM $\phi$  versus 161.4 for AM $\phi$ ). Hence, the absence of PS-R' expression by AM $\phi$  did not appear to explain our previous finding of decreased phagocytosis of apoptotic cells (10).

To determine whether PS-R' was functional on both types of M $\phi$ , we performed phagocytosis and adhesion assays in the presence of the blocking mAb 217. Blockade of PS-R' specifically and profoundly inhibited phagocytosis of apoptotic thymocytes (Fig. 9, A and B), in agreement with previous data using this mAb on other cell types (12). However, the same concentration of mAb 217 had no effect on adhesion of apoptotic thymocytes to either type of M $\phi$  (Fig. 9, C and D). These data confirm that both AM $\phi$  and PM $\phi$  express functional PS-R' but that this receptor does not mediate apoptotic cell adhesion.

To determine whether stimulation via PS-R' induced PKC activation, we tested whether mAb 217 could inhibit PS liposome-induced translocation of PKC  $\beta$ II to particulate fractions. In the presence of control IgM, PS liposomes induced translocation of PKC  $\beta$ II from cytosolic fractions to membrane and cytoskeletal fractions (Fig. 10, top row), as anticipated from our previous results (Fig. 7). Such induced translocation was inhibited when PS-R' was blocked using mAb 217 (Fig. 10, bottom row). These results indicate that the induction of PKC  $\beta$ II translocation by PS liposomes and, by implication, by apoptotic cells is mediated through PS-R'. Because mAb 217 itself has been shown to have agonist activity as indicated by its ability to stimulate transforming growth factor- $\beta$  expression (12), in separate experiments, we tested whether this mAb in the absence of PS liposomes could induce PKC translocation. No induction of PKC translocation was observed for PKC  $\beta$ II or any of the other seven isoforms tested.<sup>2</sup> Thus, the effects shown in Fig. 10 result from inhibition of PS-induced translocation mediated via PS-R' and not from agonist activity of mAb 217 itself.



**FIG. 8. Resident murine AM $\phi$  and PM $\phi$  express PS-R'.** A, reverse transcriptase-PCR analysis of mRNA. Equally loaded PCRs were normalized to achieve the same expression of the housekeeping gene glyceraldehyde-3-phosphate dehydrogenase (right side) by adjusting the volume of the cDNA product used in each reaction. The black and white image was inverted after scanning. Neg is a negative control, where no reverse transcriptase reaction was performed and aliquots of RNAs from both resident M $\phi$  were mixed. The reference ladder is 1 kb Plus (Invitrogen). Results are representative of two experiments performed with identical outcomes. B and C, flow cytometric analysis of surface protein expression. AM $\phi$  and PM $\phi$  from normal C57BL/6 mice were stained in suspension with mAb 217 culture supernatant or with control IgM from a murine myeloma, with biotinylated goat anti-mouse IgM F(ab')<sub>2</sub>, and with streptavidin/phycoerythrin and then analyzed by flow cytometry, counting 10,000 cells per condition. B, AM $\phi$ ; C, PM $\phi$ . The narrow line depicts staining with isotype control, and the dark line depicts staining with mAb 217. Representative histograms are shown; similar results were obtained in four independent experiments.

## DISCUSSION

The principal findings of this study indicate that PKC  $\beta$ II is required for phagocytosis of apoptotic cells by murine tissue M $\phi$  and that the activation of this PKC isozyme in response to PS liposomes is mediated by PS-R'. These conclusions are based on a consistent body of evidence, including the effects on phagocytosis of PKC depletion by overnight PMA treatment, dose-specific inhibition using rottlerin and Ro-32-0432, and a specific myristoylated blocking peptide corresponding to the carboxyl terminus of PKC  $\beta$ II as well as by inhibition of PS-induced translocation of PKC  $\beta$ II when PS-R' was blocked. Because adhesion of apoptotic thymocytes to M $\phi$  was not decreased by the PKC inhibitor staurosporine (11) or by overnight PMA treatment, the inhibition of phagocytosis seen in this study appears to result from recognition events following binding. Our results also showed that exposure to PS liposomes induced translocation of several other PKC isoforms ( $\beta$ I,  $\delta$ ,  $\epsilon$ ,  $\mu$ , and  $\zeta$ , and to a lesser degree  $\alpha$ ) to membrane and cytoskeletal fractions. However, based on results of the inhibitor and PMA studies, translocation of these latter isoforms appears to be unnecessary for phagocytosis. Furthermore, our results indicated that the stereo-specific PS-R' did not mediate adhesion of apoptotic cells, consistent with its predicted short extracellular domain. Finally, Western blot analysis showed that murine AM $\phi$  have markedly lower expression of PKC  $\beta$ II and of multiple other PKC isoforms ( $\alpha$ ,  $\beta$ I,  $\delta$ ,  $\epsilon$ ,  $\mu$ , and  $\zeta$ ), providing a partial explanation for the previously demonstrated relative deficit in apoptotic cell phagocytosis by that cell type (10). These novel findings advance the understanding of signal transduction during M $\phi$  recognition and ingestion of apoptotic cells.




FIG. 9. mAb against PS-R' blocks Mφ phagocytosis but not adhesion of apoptotic thymocytes. Resident AMφ and PMφ from normal C57BL/6 mice were pretreated with saturating amounts of control antibody (gray bars) or anti-PS-R' mAb 217 (black bars), and then phagocytosis or adhesion of apoptotic thymocytes was determined by examining hematoxylin-eosin Y-stained slides under oil immersion. A and B, phagocytosis. Data are mean  $\pm$  S.E. of 3–5 replicates per condition in a single experiment. \*,  $p < 0.05$ , unpaired  $t$  test. C and D, adhesion. Data are mean  $\pm$  S.E. of 3–5 replicates per condition in a single experiment. Similar results were found in an additional experiment.




FIG. 10. Antibody against PS-R' inhibits PKC  $\beta$ II translocation induced by PS liposomes. Resident PMφ were incubated with saturating amounts of control rabbit IgM (top row) or anti-PS-R' mAb 217 (bottom row) and exposed to PS liposomes to a final concentration of 0.11 mM for up to 15 min at 37 °C and 5% CO<sub>2</sub>, and then PKC  $\beta$ II expression in cytosolic versus membrane and cytoskeletal fractions was assayed by Western analysis. C, control (no liposomes).

The finding that PKC  $\beta$ II is necessary for Mφ phagocytosis of apoptotic cells is important because it differs from previously defined requisites for Mφ phagocytosis of other types of particles. Allen and Aderem (49) found that during ingestion of zymosan by lipopolysaccharide-stimulated PMφ, PKC  $\alpha$  and myristoylated, alanine-rich protein kinase C substrate co-localized with F-actin and talin adjacent to nascent phagocytic cups. During FcγR-mediated phagocytosis, PKC  $\alpha$  (50),  $\beta$  (51),  $\gamma$  (52),  $\delta$  (53), and  $\epsilon$  (50) have all been shown to localize to the phagosome membrane, with the specific PKC isoform recruited dependent on the state of Mφ differentiation and the exact FcγR involved (52). However, because PKC co-localization may be a consequence rather than a necessary process, other groups have investigated blockade of PKC function. Overexpression of a dominant negative mutant of PKC  $\alpha$  in the murine Mφ cell line RAW 264.7 reduced phagocytosis of IgG-opsonized sheep red blood cells (54) but not phagocytosis of *Leishmania donovani* promastigotes (55). Although the former finding was interpreted to indicate a requirement for PKC  $\alpha$  in FcγR-mediated phagocytosis, such observations must be interpreted with caution, because overexpression of one PKC isozyme can alter the levels of other PKC isoforms (56). By contrast, using a combination of confocal microscopy, various inhibitors, and biochemical evidence of PKC translocation in RAW 264.7 cells, Larsen *et al.* (50) found that PKC  $\alpha$  was needed for FcγR-

stimulated respiratory burst but that only PKC  $\delta$  and  $\epsilon$  were necessary for FcγR-mediated phagocytosis itself. PKC  $\beta$ II and its anchoring protein RACK1 have been found to be up-regulated in rat AMφ upon maturation of functional responses such as tumor necrosis factor- $\alpha$  or hydrogen peroxide production and lysozyme release (57). Significantly, however, none of these previous studies have confirmed a role for PKC  $\beta$ II in Mφ phagocytosis of other particles.

PKC  $\beta$ I and  $\beta$ II are products of alternative mRNA splicing of a single gene; they are identical for the first 621 amino acids and differ only in their carboxyl-terminal 50–52 amino acids. The myristoylated blocking peptides we used correspond to this area of difference and therefore were specific. The absence of effect of myristoylated  $\beta$ I peptide on phagocytosis agrees with previous reports that these isoforms have unique functions (28). Early studies found that although both isoforms translocate to membranes in response to short term PMA exposure, only PKC  $\beta$ II is an actin-binding protein that translocates to the actin-based cytoskeleton (58). However, recent studies have reached conflicting conclusions, showing either that PKC  $\beta$ II does not co-distribute with actin-based cytoskeleton upon PMA treatment (59) or that PKC  $\beta$ I as well as other PKC isoforms bind to and are activated by F-actin (60, 61).

That phagocytosis of apoptotic targets might involve unique signal transduction elements should not be surprising. In contrast to the proinflammatory responses activated by phagocytosis via other pathways (1), phagocytosis of apoptotic cells is continuous during development and antiphlogistic during resolving inflammation (62–65). The finding that murine thioglycollate-elicited PMφ secreted MIP-2/CXCL2 upon ingestion of apoptotic T cells has been interpreted as a contradiction to this principle (66), but an alternative view would be that production of this chemokine in the absence of other proinflammatory signals simply recruits Mφ as part of the beneficial clean-up process. Ingestion of apoptotic T cells has been found to compromise Mφ host defense functions *in vivo* (67). Hence, we speculate that the relatively deficient phagocytosis of apoptotic leukocytes by AMφ may be a beneficial evolutionary adaptation. Even moderate suppression of chemokine and monokine secretion in the lungs may tip the balance in favor of virulent pathogens such as enteric Gram-negative bacteria and *Staphylococcus aureus* or may be sufficient to permit evasion of innate immunity by less virulent pathogens such as anaerobes.

Our current data showing that both murine AMφ and PMφ respond to PMA agrees with the preponderance of results in other systems. Heale and Speert (68) found that PMA can reverse the inability of murine AMφ to ingest *Pseudomonas aeruginosa*. Although Peters-Golden *et al.* (69) initially reported that PMA caused release of arachidonic acid in a PKC-dependent manner in rat PMφ but not AMφ, a more recent paper from that group showed that the augmentation of FcγR-mediated phagocytosis in rat AMφ by leukotrienes is PKC-dependent, indicating that AMφ can respond to PKC-stimulatory molecules (70).

Our analyses of PS-R' advance the understanding of this receptor's role in the process of apoptotic cell clearance. The gene for this receptor has been conserved with exceedingly high fidelity across greater than 600 million years of evolution between organisms as disparate as *Caenorhabditis elegans*, *Drosophila melanogaster*, and mammals (12). This conservation and the fact that it was not previously identified by genetic analysis of *C. elegans* mutants (which suggests that its deletion may be lethal) speaks to its fundamental importance. Because we show that mAb 217 does not block adhesion, the PS-R' function must depend on initial adhesion via other receptors. Thus, it is a recognition molecule that imparts specificity to the



interaction. We cannot at present exclude the possibility that only those AM $\phi$  that express high amounts of surface PS-R' participate in phagocytosis. However, the fact that only roughly 40% of PM $\phi$  showed significant staining, yet virtually all PM $\phi$  ingest apoptotic cells in an annexin V-inhibitable fashion (10), urges caution in interpretation of flow cytometry data using mAb 217. The high background staining seen using this IgM mAb precludes accurate definition of the lower limit of expression, and even small amounts of PS-R' may be sufficient. In addition to the stereo-specific PS-R', PS can be bound by multiple receptors including class B scavenger receptors, CD14, macrophage (CD68), and thrombospondin-dependent vitronectin receptors (1, 71–73).

The demonstration that resident murine AM $\phi$  express lesser amounts of multiple PKC isoforms, compared with resident M $\phi$  at another body surface, further highlights the distinctive differentiation of this cell type (74, 75). AM $\phi$  appear to have evolved as specialized phagocytes whose principal task is to keep the gas-exchanging surface of the alveoli clear yet not compromised by excessive inflammation. AM $\phi$  have a mostly suppressive role on the induction of adaptive immune responses within the lung itself (76, 77), permitting local expression of T cell effector functions while inhibiting their proliferation (78). Indeed, AM $\phi$  differ from other M $\phi$  types in expression of surface receptors (79, 80), in antigen-presenting capacity (81), and in eicosanoid production (82, 83). Our results agree with previous studies that found markedly lower expression of multiple PKC isoforms, including PKC  $\beta$ II, in normal human or rat AM $\phi$  compared with monocytes or other M $\phi$  (37, 69). Interestingly, we have also found that, like murine AM $\phi$ , the murine M $\phi$  cell lines RAW 264.7 and IC21 have markedly reduced expression of PKC  $\beta$ II (but not PKC  $\alpha$  or  $\beta$ I), which correlates with their greatly reduced phagocytosis (but not adhesion) of apoptotic thymocytes relative to PM $\phi$ .<sup>2</sup> Significantly, the fact that Monick *et al.* (37) found the greatest difference in PKC isoform expression between human AM $\phi$  and blood monocytes to be a reduction in PKC  $\beta$ II in AM $\phi$  implies that our findings can be extrapolated to normal human AM $\phi$ , which we have recently found also have relatively depressed phagocytosis of apoptotic lymphocytes.<sup>2</sup>

In summary, these data demonstrate for the first time that M $\phi$  phagocytosis of apoptotic cells requires PKC  $\beta$ II, an isoform not shown previously to be required for phagocytosis of other types of particles. This unique PKC isoform requirement may be an important clue to the signaling pathways that mediate suppression of M $\phi$  production of proinflammatory mediators on ingestion of apoptotic cells (64, 65). Activation of PKC  $\beta$ II by PS-R' provides a partial explanation for the essential role of this receptor in apoptotic cell ingestion.

**Acknowledgments**—We thank Dr. Lilli Petruzzelli and all of the members of the Ann Arbor Veterans Affairs Medical Center Research Enhancement Award Program for helpful suggestions, Dr. Valerie Fadok for the generous gift of mAb 217, and Joyce O'Brien for secretarial support.

# REFERENCES

1. Aderem, A., and Underhill, D. M. (1999) *Annu. Rev. Immunol.* **17**, 593–623
2. Fadok, V. A., and Chimini, G. (2001) *Semin. Immunol.* **13**, 365–372
3. Savill, J. S., Wyllie, A. H., Henson, J. E., Walport, M. J., Henson, P. M., and Haslett, C. (1989) *J. Clin. Invest.* **83**, 865–875
4. Akbar, A. N., Savill, J., Gombert, W., Boffill, M., Borthwick, N. J., Whitelaw, F., Grundy, J., Janossy, G., and Salmon, M. (1994) *J. Exp. Med.* **180**, 1943–1947
5. Stern, M., Savill, J., and Haslett, C. (1996) *Am. J. Pathol.* **149**, 911–921
6. Haslett, C. (1999) *Am. J. Respir. Crit. Care Med.* **160**, S5–S11
7. Surh, C. D., and Sprent, J. (1994) *Nature* **372**, 100–103
8. Milik, A. M., Buechner-Maxwell, V. A., Sonstein, J., Kim, S., Seitzman, G. D., Beals, T. F., and Curtis, J. L. (1997) *J. Clin. Invest.* **99**, 1082–1091
9. Newman, S. L., Henson, J. E., and Henson, P. M. (1982) *J. Exp. Med.* **156**, 430–442
10. Hu, B., Sonstein, J., Christensen, P. J., Punturieri, A., and Curtis, J. L. (2000) *J. Immunol.* **165**, 2124–2133

11. Hu, B., Punturieri, A., Todt, J., Sonstein, J., Polak, T., and Curtis, J. L. (2002) *J. Leukocyte Biol.* **71**, 881–889
12. Fadok, V. A., Bratton, D. L., Rose, D. M., Pearson, A., Ezekewitz, R. A., and Henson, P. M. (2000) *Nature* **405**, 85–90
13. Verhoven, B., Krahling, S., Schlegel, R. A., and Williamson, P. (1999) *Cell Death Differ.* **6**, 262–270
14. Schlegel, R. A., Stevens, M., Lumley-Sapanski, K., and Williamson, P. (1993) *Immunol. Lett.* **36**, 283–288
15. Koopman, G., Reutelingsperger, C. P., Kuijten, G. A., Keehnen, R. M., Pals, S. T., and van Oers, M. H. (1994) *Blood* **84**, 1415–1420
16. Homburg, C. H., de Haas, M., von dem Borne, A. E., Verhoeven, A. J., Reutelingsperger, C. P., and Roos, D. (1995) *Blood* **85**, 532–540
17. Martin, S. J., Reutelingsperger, C. P., McGahon, A. J., Rader, J. A., van Schie, R. C., LaFace, D. M., and Green, D. R. (1995) *J. Exp. Med.* **182**, 1545–1556
18. Fadok, V. A., de Cathelineau, A., Daleke, D. L., Henson, P. M., and Bratton, D. L. (2001) *J. Biol. Chem.* **276**, 1071–1077
19. Scott, R. S., McMahon, E. J., Pop, S. M., Reap, E. A., Caricchio, R., Cohen, P. L., Earp, H. S., and Matsushima, G. K. (2001) *Nature* **411**, 207–211
20. Gregory, C. D. (2000) *Curr. Opin. Immunol.* **12**, 27–34
21. Hoffmann, P. R., deCathelineau, A. M., Ogden, C. A., Leverrier, Y., Bratton, D. L., Daleke, D. L., Ridley, A. J., Fadok, V. A., and Henson, P. M. (2001) *J. Cell Biol.* **155**, 649–659
22. Platt, N., da Silva, R. P., and Gordon, S. (1999) *Immunol. Lett.* **65**, 15–19
23. Marguet, D., Luciani, M. F., Moynault, A., Williamson, P., and Chimini, G. (1999) *Nat. Cell Biol.* **1**, 454–456
24. Webb, B. L., Hirst, S. J., and Giembycz, M. A. (2000) *Br. J. Pharmacol.* **130**, 1433–1452
25. Newton, A. C. (1995) *J. Biol. Chem.* **270**, 28495–28498
26. Mochly-Rosen, D., and Gordon, A. S. (1998) *FASEB J.* **12**, 35–42
27. Dempsey, E. C., Newton, A. C., Mochly-Rosen, D., Fields, A. P., Reyland, M. E., Insel, P. A., and Messing, R. O. (2000) *Am. J. Physiol.* **279**, L429–L438
28. Chalfant, C. E., Ohno, S., Konno, Y., Fisher, A. A., Bisnauth, L. D., Watson, J. E., and Cooper, D. R. (1996) *Mol. Endocrinol.* **10**, 1273–1281
29. Gokmen-Polar, Y., and Fields, A. P. (1998) *J. Biol. Chem.* **273**, 20261–20266
30. Martiny-Baron, G., Kazanietz, M. G., Mischak, H., Blumberg, P. M., Kochs, G., Hug, H., Marme, D., and Schachtele, C. (1993) *J. Biol. Chem.* **268**, 9194–9197
31. Keenan, C., Goode, N., and Pears, C. (1997) *FEBS Lett.* **415**, 101–108
32. Way, K. J., Chou, E., and King, G. L. (2000) *Trends Pharmacol. Sci.* **21**, 181–187
33. Curtis, J. L., Kim, S., Scott, P. J., and Buechner-Maxwell, V. A. (1995) *Am. J. Respir. Cell Mol. Biol.* **12**, 520–530
34. Stebbins, E. G., and Mochly-Rosen, D. (2001) *J. Biol. Chem.* **276**, 29644–29650
35. Zar, J. H. (1974) *Biostatistical Analysis*, 2nd Ed., pp. 122–165, Prentice-Hall, Englewood Cliffs, NJ
36. Licht, R., Jacobs, C. W., Tax, W. J., and Berden, J. H. (1999) *J. Immunol. Methods* **223**, 237–248
37. Monick, M. L., Carter, A. B., Gudmundsson, G., Geist, L. J., and Hunninghake, G. W. (1998) *Am. J. Physiol.* **275**, L389–L397
38. Nixon, J. B., and McPhail, L. C. (1999) *J. Immunol.* **163**, 4574–4582
39. Ahnadi, C. E., Giguere, P., Gravel, S., Gagne, D., Goulet, A. C., Fulop, T., Jr., Payet, M. D., and Dupuis, G. (2000) *J. Leukocyte Biol.* **68**, 293–300
40. Gschwendt, M., Muller, H. J., Kielbassa, K., Zang, R., Kittstein, W., Rincke, G., and Marks, F. (1994) *Biochem. Biophys. Res. Commun.* **199**, 93–98
41. Wilkinson, S. E., Parker, P. J., and Nixon, J. S. (1993) *Biochem. J.* **294**, 335–337
42. Keranen, L. M., and Newton, A. C. (1997) *J. Biol. Chem.* **272**, 25959–25967
43. Staudinger, J., Lu, J., and Olson, E. N. (1997) *J. Biol. Chem.* **272**, 32019–32024
44. Mochly-Rosen, D., and Kauvar, L. M. (2000) *Semin. Immunol.* **12**, 55–61
45. Fadok, V. A., Savill, J. S., Haslett, C., Bratton, D. L., Doherty, D. E., Campbell, P. A., and Henson, P. M. (1992) *J. Immunol.* **149**, 4029–4035
46. Tait, J. F., and Smith, C. (1999) *J. Biol. Chem.* **274**, 3048–3054
47. Raz, A., Bucana, C., Fogler, W. E., Poste, G., and Fidler, I. J. (1981) *Cancer Res.* **41**, 487–494
48. Perry, D. G., and Martin, W. J., II (1995) *J. Immunol. Methods* **181**, 269–285
49. Allen, L. H., and Aderem, A. (1995) *J. Exp. Med.* **182**, 829–840
50. Larsen, E. C., DiGennaro, J. A., Saito, N., Mehta, S., Loegering, D. J., Mazurkiewicz, J. E., and Lennartz, M. R. (2000) *J. Immunol.* **165**, 2809–2817
51. Dekker, L. V., Leitges, M., Altschuler, G., Mistry, N., McDermott, A., Roes, J., and Segal, A. W. (2000) *Biochem. J.* **347**, 285–289
52. Melendez, A. J., Harnett, M. M., and Allen, J. M. (1999) *Immunology* **96**, 457–464
53. Brumell, J. H., Howard, J. C., Craig, K., Grinstein, S., Schreiber, A. D., and Tyers, M. (1999) *J. Immunol.* **163**, 3388–3395
54. Breton, A., and Descoteaux, A. (2000) *Biochem. Biophys. Res. Commun.* **276**, 472–476
55. St-Denis, A., Caouras, V., Gervais, F., and Descoteaux, A. (1999) *J. Immunol.* **163**, 5505–5511
56. Ways, D. K., Kukoly, C. A., deVente, J., Hooker, J. L., Bryant, W. O., Posekany, K. J., Fletcher, D. J., Cook, P. P., and Parker, P. J. (1995) *J. Clin. Invest.* **95**, 1906–1915
57. Corsini, E., Viviani, B., Lucchi, L., Marinovich, M., Racchi, M., and Galli, C. L. (2001) *Immunol. Lett.* **76**, 89–93
58. Blobe, G. C., Stribling, D. S., Fabbro, D., Stabel, S., and Hannun, Y. A. (1996) *J. Biol. Chem.* **271**, 15823–15830
59. Edwards, A. S., Faux, M. C., Scott, J. D., and Newton, A. C. (1999) *J. Biol. Chem.* **274**, 6461–6468
60. Slater, S. J., Milano, S. K., Stagliano, B. A., Gergich, K. J., Curry, J. P., Taddeo, F. J., and Stubbs, C. D. (2000) *Biochemistry* **39**, 271–280
61. Lopez-Lluch, G., Bird, M. M., Canas, B., Godovac-Zimmerman, J., Ridley, A., Segal, A. W., and Dekker, L. V. (2001) *Biochem. J.* **357**, 39–47



62. Meagher, L. C., Savill, J. S., Baker, A., Fuller, R. W., and Haslett, C. (1992) *J. Leukocyte Biol.* **52**, 269–273
63. Voll, R. E., Herrmann, M., Roth, E. A., Stach, C., Kalden, J. R., and Girkontaite, I. (1997) *Nature* **390**, 350–351
64. Fadok, V. A., Bratton, D. L., Konowal, A., Freed, P. W., Westcott, J. Y., and Henson, P. M. (1998) *J. Clin. Invest.* **101**, 890–898
65. McDonald, P. P., Fadok, V. A., Bratton, D., and Henson, P. M. (1999) *J. Immunol.* **163**, 6164–6172
66. Uchimura, E., Kodaira, T., Kurosaka, K., Yang, D., Watanabe, N., and Kobayashi, Y. (1997) *Biochem. Biophys. Res. Commun.* **239**, 799–803
67. Freire-de-Lima, C. G., Nascimento, D. O., Soares, M. B. P., Bozza, P. T., Castro-Faria-Neto, H. C., de Mello, F. G., DosReis, G. A., and Lopes, M. F. (2000) *Nature* **403**, 199–203
68. Heale, J. P., and Speert, D. P. (2001) *J. Leukocyte Biol.* **69**, 158–160
69. Peters-Golden, M., McNish, R. W., Sporn, P. H., and Balazovich, K. (1991) *Am. J. Physiol.* **261**, L462–L471
70. Mancuso, P., and Peters-Golden, M. (2000) *Am. J. Respir. Cell Mol. Biol.* **23**, 727–733
71. Savill, J., Fadok, V., Henson, P., and Haslett, C. (1993) *Immunol. Today* **14**, 131–136
72. Devitt, A., Moffatt, O. D., Raykundalia, C., Capra, J. D., Simmons, D. L., and Gregory, C. D. (1998) *Nature* **392**, 505–509
73. Ramprasad, M. P., Fischer, W., Witztum, J. L., Sambrano, G. R., Quehenberger, O., and Steinberg, D. (1995) *Proc. Natl. Acad. Sci. U. S. A.* **92**, 9580–9584
74. Lipscomb, M. F., and Russell, S. W. (eds) (1997) *Lung Macrophages and Dendritic Cells in Health and Disease*, Vol. 107 (Lenfant, C., ed) Marcel Dekker, New York
75. Brody, A. R. (1998) *J. Lab. Clin. Med.* **131**, 391–392
76. Thepen, T., Van Rooijen, N., and Kraal, G. (1989) *J. Exp. Med.* **170**, 499–509
77. Holt, P. G., Oliver, J., Bilyk, N., McMenamin, C., McMenamin, P. G., Kraal, G., and Thepen, T. (1993) *J. Exp. Med.* **177**, 397–407
78. Upham, J. W., Strickland, D. H., Bilyk, N., Robinson, B. W. S., and Holt, P. G. (1995) *Immunology* **84**, 142–147
79. Hance, A. J., Douches, S., Winchester, R. J., Ferrans, V. J., and Crystal, R. G. (1985) *J. Immunol.* **134**, 284–292
80. Yoshikawa, K., Suzuki, Y., Kawai, M., Fukada, M., and Yokochi, T. (1991) *Microbiol. Immunol.* **35**, 803–807
81. Lipscomb, M. F., Lyons, C. R., Nunez, G., Ball, E. J., Stanstny, P., Vial, W., Lem, V., Weissler, J. C., Miller, L. M., and Toews, G. B. (1986) *J. Immunol.* **136**, 497–504
82. Bigby, T. D., and Holtzman, M. J. (1987) *J. Immunol.* **138**, 1546–1550
83. Balter, M. S., Toews, G. B., and Peters-Golden, M. (1989) *J. Immunol.* **142**, 602–608

**Activation of Protein Kinase C  $\beta$ II by the Stereo-specific Phosphatidylserine Receptor Is Required for Phagocytosis of Apoptotic Thymocytes by Resident Murine Tissue Macrophages**

Jill C. Todt, Bin Hu, Antonello Punturieri, Joanne Sonstein, Timothy Polak and Jeffrey L. Curtis

*J. Biol. Chem.* 2002, 277:35906-35914.

doi: 10.1074/jbc.M202967200 originally published online July 11, 2002

---

Access the most updated version of this article at doi: [10.1074/jbc.M202967200](https://doi.org/10.1074/jbc.M202967200)

Alerts:

- [When this article is cited](#)
- [When a correction for this article is posted](#)

[Click here](#) to choose from all of JBC's e-mail alerts

This article cites 81 references, 34 of which can be accessed free at <http://www.jbc.org/content/277/39/35906.full.html#ref-list-1>