

Histidine 129 in the 75-kDa Subunit of Mitochondrial Complex I from *Yarrowia lipolytica* Is Not a Ligand for [Fe₄S₄] Cluster N5 but Is Required for Catalytic Activity*

Received for publication, October 8, 2004, and in revised form, November 24, 2004
Published, JBC Papers in Press, November 30, 2004, DOI 10.1074/jbc.M411488200

Antje Waletko, Klaus Zwicker, Albina Abdrakhmanova, Volker Zickermann, Ulrich Brandt, and Stefan Kerscher‡

From the Universität Frankfurt, Fachbereich Medizin, Institut für Biochemie I, F-60590 Frankfurt am Main, Germany

Respiratory chain complex I contains 8–9 iron-sulfur clusters. In several cases, the assignment of these clusters to subunits and binding motifs is still ambiguous. To test the proposed ligation of the tetranuclear iron-sulfur cluster N5 of respiratory chain complex I, we replaced the conserved histidine 129 in the 75-kDa subunit from *Yarrowia lipolytica* with alanine. In the mutant strain, reduced amounts of fully assembled but destabilized complex I could be detected. Deamino-NADH: ubiquinone oxidoreductase activity was abolished completely by the mutation. However, EPR spectroscopic analysis of mutant complex I exhibited an unchanged cluster N5 signal, excluding histidine 129 as a cluster N5 ligand.

Respiratory chain complex I links the transfer of two electrons from NADH to ubiquinone to the translocation of four protons across the respiratory membrane (1). Only low resolution structures of complex I are available, demonstrating that the mitochondrial enzyme is L-shaped and consists of a membrane arm and a peripheral arm (2–4). The reaction mechanism is largely unknown. Electrons from NADH are first transferred to FMN, non-covalently bound to the 51-kDa subunit, then to a series of iron-sulfur clusters in the peripheral arm, and finally to the ubiquinone reduction site which, as has been demonstrated by site-directed mutagenesis in the obligate aerobic yeast *Yarrowia lipolytica*, is partly contained in the 49-kDa subunit of the peripheral arm (5). Localization by electron microscopic single particle analysis of two anti-49-kDa subunit monoclonal antibody epitopes has provided evidence that this catalytic core of complex I is located ~70–80 Å away from the phospholipid bilayer (6). From these and other results, an indirect coupling mechanism between quinone reduction in the peripheral arm and proton pumping in the membrane arm seems likely (1, 7, 8).

Little information is available on the spatial arrangement of the other components of the electron transfer pathway. EPR¹-detectable iron-sulfur clusters have been assigned to individual subunits (9) as follows: the 75-kDa subunit, containing the [Fe₂S₂] cluster N1b and the [Fe₄S₄] clusters N4 and N5; the 51-kDa subunit, containing [Fe₄S₄] cluster N3; the 24-kDa

subunit, containing the [Fe₂S₂] cluster N1a; and the PSST subunit, containing the [Fe₄S₄] cluster N2. Two EPR-silent [Fe₄S₄] clusters, termed N6a and N6b, have been assigned to the ferredoxin-like subunit of *Neurospora crassa* complex I homologous to bovine TYKY (10). The tetranuclear iron-sulfur cluster N5 is only observed at very low temperatures and high microwave power. After its initial discovery in pigeon heart complex I (11), it has also been identified in bovine (9), *Rhodobacter spaeroides* (12), and *Y. lipolytica* (4, 13), but not in *N. crassa* (14) complex I. In *Y. lipolytica*, cluster N5 shows very fast spin relaxation with a half-saturation parameter of ~180 milliwatts at 5 K. The line width of the g_z signal (L_z) is in the range of 2.5–3 millitesla. Similar to observations in other model systems, the spin concentration of cluster N5 is much lower than that of the other EPR-detectable clusters (13).

The N-terminal part of the 75-kDa subunit contains three highly conserved iron-sulfur cluster motifs and is homologous to the N-terminal portion of iron-only hydrogenases (15). In the x-ray structure of the iron-only hydrogenase from *Clostridium pasteurianum*, these motifs ligate one [Fe₂S₂] and two [Fe₄S₄] iron-sulfur clusters (16). Recently, direct evidence for the presence of three iron-sulfur clusters in the heterologously expressed 75-kDa subunit from *Paracoccus denitrificans* has been presented (17). In the same study, the fast relaxing [Fe₄S₄] cluster N5 has been assigned to the second binding motif of the 75-kDa subunit, an unusual HXXXXCXXCXXXXXC motif that contains three cysteines and a histidine. Secondary structure predictions using the PROF algorithm (26) suggest that this sequence element, both in hydrogenases and complex I from various organisms, forms a loop bounded by α-helices (Fig. 1). To test the proposed assignment of this sequence as a cluster N5 ligation motif in fully assembled complex I, we mutated the conserved histidine 129 of the *Y. lipolytica* 75-kDa subunit to alanine.

MATERIALS AND METHODS

The *Y. lipolytica* deletion strain *nuamΔ* (*ura3-302, leu2-270, lys11-23, nuam::URA3, NUGM-Htg2, NDH2i, MatB*), in which a 1.7-kb *Cl*I/*K*p*n*I fragment corresponding to codons 34–598 of the *NUAM* gene encoding the 728-amino acid precursor of the 75-kDa subunit of complex I is replaced with the *Y. lipolytica URA3* gene (1.6 kb) oriented in opposite direction to the original *NUAM* open reading frame, was constructed by the double homologous recombination strategy published previously (18). A wild type and a PCR-generated H129A mutant version of the *NUAM* gene were then subcloned as 5.2-kb *S*alI fragments (after an internal *S*alI site in the *NUAM* open reading frame had been mutagenized to GTTGAC) into the replicative plasmid pUB4 (19) and transformed into strain *nuamΔ*, resulting in the strains *nuamΔ*, p*NUAM*-wild type (parental) and *nuamΔ*, p*NUAM*-H129A (mutant). The presence of the point mutation in large scale cultures of strain *nuamΔ*, p*NUAM*-H129A was verified by the isolation of total DNA from 5-ml aliquots and the direct sequencing of PCR products.

Complex I catalytic activities in *Y. lipolytica* mitochondrial mem-

* This work was supported by a grant from the Deutsche Forschungsgemeinschaft for Sonderforschungsbereich 472, project P2. The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked "advertisement" in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

‡ To whom correspondence should be addressed. Tel.: 49-6301-6943; Fax: 49-6301-6970; E-mail: kerscher@zbc.kgu.de.

¹ The abbreviations used are: EPR, electron paramagnetic resonance; BN, blue native; HAR, hexa-ammine-ruthenium(III)-chloride.

TABLE I
NADH:HAR and dNADH:DBQ activities of mitochondrial membranes

The abbreviations used here that are not defined elsewhere are as follows: DBQ, n-decylubiquinone; dNADH, deamino-NADH; WT, wild type.

Strain	NADH:HAR activity		dNADH:DBQ activity	
	$\mu\text{mol min}^{-1} \text{mg}^{-1}$	Percent of activity	$\mu\text{mol min}^{-1} \text{mg}^{-1}$	Percent of activity
<i>Nuam</i> Δ ,pNUAM-WT	0.84	100	0.290	100
<i>Nuam</i> Δ ,pNUAM-H129A	0.62	74	<0.01	<3
<i>Nuam</i> Δ ,pUB4	0.28	33	<0.01	<3




FIG. 3. EPR spectra of purified complex I from mutant H129A (6.3 mg/ml) and parental strain (3.7 mg/ml). A microwave frequency of 9.47 GHz and a modulation amplitude of 0.64 millitesla were used, as well as a spectrum at 40 K and a microwave power of 1 milliwatt (A), a spectrum at 12 K and a microwave power of 1 milliwatt (B), and a spectrum at 5 K and a microwave power of 100 milliwatt (C). Characteristic signal positions of individual clusters are indicated. The region between $g = 1.9$ and $g = 1.95$ reflects contributions from all iron-sulfur clusters detectable under these conditions.

signature of cluster N5 was completely unchanged in complex I purified from the mutant strain, we could exclude this possibility. Our results are in contradiction with a recent study with a heterologously expressed 75-kDa subunit from *P. denitrificans* (17). From changes in the EPR spectrum of the iron-sulfur cluster N5 that were caused by a mutation changing the corresponding histidine to a cysteine, the authors concluded that the HXXXXCXXXXXXC motif of the 75-kDa subunit binds cluster N5. The most obvious explanation for the results obtained with the *P. denitrificans* H106C mutant would be that

structural changes had indirectly changed the geometry of cluster N5. Such EPR signal reduction, peak broadening, and shifting effects have been described, e.g. for point mutations affecting amino acids in the vicinity of cluster N2 of complex I from *Y. lipolytica* (5, 24, 25), and can be expected to be more pronounced when single subunits instead of completely assembled multi-subunit complexes are studied. In fact, significant instability of the mutant *P. denitrificans* 75-kDa subunit was noted (17). Although we also observed destabilization of *Y. lipolytica* complex I by the H129A mutation, it obviously did not affect the EPR spectra of the purified enzyme.

A more fundamental problem with complex I from *P. denitrificans* is that the g signal values of cluster N5, detected at 5 K, are very close to those of cluster N4, detected at 12 K (17). Distinction between these clusters relies on the assumption that the cluster N4 signals of *P. denitrificans* complex I are completely power saturated at 5 K and a microwave power of 100 milliwatt, which in our hands is clearly not the case for cluster N4 of *Bos taurus* and *Y. lipolytica* complex I. Therefore, although at least the g_z signals of clusters N4 and N5 are well separated in the latter two model organisms, the assignment of EPR signals to cluster N5 of *P. denitrificans* complex I is more difficult. It should be noted that the cluster N4 EPR signals were also found to be somewhat affected by the H106C mutation in *P. denitrificans* complex I (17). Thus, there is another possibility in regard to how the findings in *P. denitrificans* and *Y. lipolytica* could be reconciled. If the iron-sulfur cluster assignment within the 75-kDa subunit was revised, the mutation of the conserved histidine in *Y. lipolytica* may have resulted in the removal of a cluster N1b ligand, leaving the EPR signature of cluster N5 unchanged. The histidine to cysteine exchange in *P. denitrificans* as such may then have been EPR-silent, but it may have caused a structural alteration that affected the EPR patterns of both clusters N4 and N5.

However, because the study by Yano *et al.* (17) clearly demonstrates the presence of three iron-sulfur clusters in the heterologously expressed 75-kDa subunit of *P. denitrificans*, this proposal raises another problem. If all three motifs in the 75-kDa hold an iron-sulfur cluster, why did the H129A mutation not have any effect on the EPR spectra of *Y. lipolytica* complex I? Possible answers are that cluster N1b is either not present or undetectable by standard EPR spectroscopy in complex I from *Y. lipolytica*. Although complex I from bovine heart mitochondria and various other sources is known to contain two EPR detectable binuclear iron-sulfur clusters called N1a and N1b (9), only one binuclear iron-sulfur cluster (called N1) could be detected in complex I from *Y. lipolytica* to date. The stoichiometry of clusters N1/N2/N3/N4 was determined as 1:1:1:1 (13). As the N1 signal of *Y. lipolytica* complex I was not detectable in a subcomplex² including the 75-kDa but lacking the 24-kDa subunit, which in bovine heart complex I contains binuclear cluster N1a, it seems tempting to speculate that also in the *Y. lipolytica* enzyme the 75-kDa subunit may carry a second binuclear cluster that has, to date, escaped detection by

² V. Zickermann, K. Zwicker, M. Bostina, M. Radermacher, and U. Brandt, manuscript in preparation.

EPR. If the HXXXXXXCXXXXXC motif ligated this EPR silent cluster, it would have been removed by the H129A mutation, offering a straightforward explanation of why ubiquinone reductase activity was lost.

As a consequence, one has to conclude that in complex I the use of the iron-sulfur binding motifs has changed from iron-only hydrogenases. Either the histidine in question that is conserved in all known complex I sequences has acquired a different function in complex I, or the motif ligating a binuclear cluster in hydrogenase binds a tetranuclear cluster (N5) in complex I and *vice versa*.

Acknowledgments—We thank Gudrun Beyer for excellent technical assistance and Nikki Bolcevic for help in the isolation of strain *nuam*Δ. We thank Maja Aleksandra Tocilescu for preparing the BN-PAGE gels shown in Fig. 2 and Hermann Schagger for help with interpretation of the gels.

REFERENCES

1. Brandt, U., Kerscher, S., Dröse, S., Zwicker, K., and Zickermann, V. (2003) *FEBS Lett.* **545**, 9–17
2. Guenebaut, V., Schlitt, A., Weiss, H., Leonard, K., and Friedrich, T. (1998) *J. Mol. Biol.* **276**, 105–112
3. Grigorieff, N. (1998) *J. Mol. Biol.* **277**, 1033–1046
4. Djafarzadeh, R., Kerscher, S., Zwicker, K., Radermacher, M., Lindahl, M., Schagger, H., and Brandt, U. (2000) *Biochim. Biophys. Acta* **1459**, 230–238
5. Kashani-Poor, N., Zwicker, K., Kerscher, S., and Brandt, U. (2001) *J. Biol. Chem.* **276**, 24082–24087
6. Zickermann, V., Bostina, M., Hunte, C., Ruiz, T., Radermacher, M., and Brandt, U. (2003) *J. Biol. Chem.* **278**, 29072–29078
7. Friedrich, T. (2001) *J. Bioenerg. Biomembr.* **33**, 169–177
8. Yagi, T., and Matsuno-Yagi, A. (2003) *Biochemistry* **42**, 2266–2274
9. Ohnishi, T. (1998) *Biochim. Biophys. Acta* **1364**, 186–206
10. Rasmussen, T., Scheide, D., Brors, B., Kintscher, L., Weiss, H., and Friedrich, T. (2001) *Biochemistry* **40**, 6124–6131
11. Ohnishi, T., Salerno, J. C., Winter, D. B., Lim, J., Yu, C.-A., Yu, L., and King, T. E. (1975) *Biochim. Biophys. Acta* **387**, 475–490
12. Sled, V. D., Friedrich, T., Leif, H., Weiss, H., Fukumori, Y., Calhoun, M. W., Gennis, R. B., Ohnishi, T., and Meinhardt, S. W. (1993) *J. Bioenerg. Biomembr.* **25**, 347–356
13. Kerscher, S., Kashani-Poor, N., Zwicker, K., Zickermann, V., and Brandt, U. (2001) *J. Bioenerg. Biomembr.* **33**, 187–196
14. Weiss, H., Friedrich, T., Hofhaus, G., and Preis, D. (1991) *Eur. J. Biochem.* **197**, 563–576
15. Smith, M. A., Finel, M., Korolik, V., and Mendz, G. L. (2000) *Arch. Microbiol.* **174**, 1–10
16. Peters, J. W., Lanzilotta, W. N., Lemon, B. J., and Seefeldt, L. C. (1998) *Science* **282**, 1853–1858
17. Yano, T., Sklar, J., Nakamaru-Ogiso, E., Yagi, T., and Ohnishi, T. (2003) *J. Biol. Chem.* **278**, 15514–15522
18. Kerscher, S., Dröse, S., Zwicker, K., Zickermann, V., and Brandt, U. (2002) *Biochim. Biophys. Acta* **1555**, 83–91
19. Kerscher, S., Eschemann, A., Okun, P. M., and Brandt, U. (2001) *J. Cell Sci.* **114**, 3915–3921
20. Kerscher, S., Bénit, P., Abdrakhmanova, A., Zwicker, K., Rais, I., Karas, M., Rustin, P., and Brandt, U. (2004) *Eur. J. Biochem.* **271**, 3588–3595
21. Schagger, H. (2003) in *Membrane Protein Purification and Crystallization: A Practical Guide* (Hunte, C., von Jagow, G., and Schagger, H., eds) pp. 105–130, Academic Press, San Diego, CA
22. Kashani-Poor, N., Kerscher, S., Zickermann, V., and Brandt, U. (2001) *Biochim. Biophys. Acta* **1504**, 363–370
23. Gavrikova, E. V., Grivennikova, V. G., Sled, V. D., Ohnishi, T., and Vinogradov, A. D. (1995) *Biochim. Biophys. Acta* **1230**, 23–30
24. Ahlers, P., Zwicker, K., Kerscher, S., and Brandt, U. (2000) *J. Biol. Chem.* **275**, 23577–23582
25. Garofano, A., Zwicker, K., Kerscher, S., Okun, P., and Brandt, U. (2003) *J. Biol. Chem.* **278**, 42435–42440
26. Ouali, M., and King, R. D. (2000) *Protein Sci.* **9**, 1162–1176

Histidine 129 in the 75-kDa Subunit of Mitochondrial Complex I from *Yarrowia lipolytica* Is Not a Ligand for [Fe₄S₄] Cluster N5 but Is Required for Catalytic Activity

Antje Waletko, Klaus Zwicker, Albina Abdrakhmanova, Volker Zickermann, Ulrich Brandt and Stefan Kerscher

J. Biol. Chem. 2005, 280:5622-5625.

doi: 10.1074/jbc.M411488200 originally published online November 30, 2004

Access the most updated version of this article at doi: [10.1074/jbc.M411488200](https://doi.org/10.1074/jbc.M411488200)

Alerts:

- [When this article is cited](#)
- [When a correction for this article is posted](#)

[Click here](#) to choose from all of JBC's e-mail alerts

This article cites 25 references, 7 of which can be accessed free at <http://www.jbc.org/content/280/7/5622.full.html#ref-list-1>