

Additions and Corrections

Vol. 279 (2004) 34240–34249

The sensitivity of G protein-activated K⁺ channels toward halothane is essentially determined by the C terminus.

Sergej Milovic, Bibiane Steinecker-Frohnwieser, Wolfgang Schreibmayer, and Lukas G. Weigl

Page 34240: An error appeared in the funding footnote. The funding footnote should read as follows:

This work was supported by Grant 8266 from the Oesterreichische Nationalbank (to L. G. W.), Austrian Research Foundation Grants T124-B04 (to B. S.-F.) and SFB007/F708 (to W. S.), and Austrian Ministry of Science Grant UGP4 (to W. S.).

We suggest that subscribers photocopy these corrections and insert the photocopies at the appropriate places where the article to be corrected originally appeared. Authors are urged to introduce these corrections into any reprints they distribute. Secondary (abstract) services are urged to carry notice of these corrections as prominently as they carried the original abstracts.

The sensitivity of G protein-activated K⁺ channels toward halothane is essentially determined by the C terminus. Vol. 279 (2004) 34240–34249

Sergej Milovic, Bibiane Steinecker-Frohnwieser, Wolfgang Schreibmayer and Lukas G. Weigl

J. Biol. Chem. 2005, 280:6252.

Access the most updated version of this article at <http://www.jbc.org/content/280/7/6252>

Alerts:

- [When this article is cited](#)
- [When a correction for this article is posted](#)

[Click here](#) to choose from all of JBC's e-mail alerts

This article cites 0 references, 0 of which can be accessed free at <http://www.jbc.org/content/280/7/6252.full.html#ref-list-1>