

VOLUME 290 (2015) PAGES 1210–1221

DOI 10.1074/jbc.A114.608232

**The Dbf4-Cdc7 kinase promotes Mcm2-7 ring opening to allow for single-stranded DNA extrusion and helicase assembly.**

Irina Bruck and Daniel L. Kaplan

This article has been withdrawn by the authors. Fig. 1, *A* and *B*, contained missing data. The phospho-Mcm2 antibody in Fig. 1*C* did not accurately represent experimental conditions. The Mcm2-His immunoblots in Fig. 4*A* were duplicated. The Sld3-Myc immunoblot from whole cell extracts in Fig. 4*A* was inappropriately manipulated. The Sld3-Myc immunoblot from the co-immunoprecipitation experiment in Fig. 4*A* was a composite. A portion of the Psf2-5Flag immunoblot from the co-immunoprecipitation experiment from Fig. 4*A* was duplicated from the Cdc45-6HA immunoblot from the same figure panel. The Cdc45-6HA immunoblot from whole cell extracts in Fig. 4*A* and the Psf2-5Flag immunoblot from the co-immunoprecipitation experiment in Fig. 4*B* were duplicated. *Lane 1* from the Sld3-Myc immunoblot from the co-immunoprecipitation experiment in Fig. 4*B* was duplicated from *lane 8* of the Sld3-Myc immunoblot from the co-immunoprecipitation experiment in Fig. 4*A*.

**The Dbf4-Cdc7 kinase promotes Mcm2-7 ring opening to allow for single-stranded DNA extrusion and helicase assembly.**

Irina Bruck and Daniel L. Kaplan

*J. Biol. Chem.* 2017, 292:10320.

doi: 10.1074/jbc.A114.608232

---

Access the most updated version of this article at <http://www.jbc.org/content/292/24/10320>

Alerts:

- [When this article is cited](#)
- [When a correction for this article is posted](#)

[Click here](#) to choose from all of JBC's e-mail alerts

This article cites 0 references, 0 of which can be accessed free at <http://www.jbc.org/content/292/24/10320.full.html#ref-list-1>