

VOLUME 285 (2010) PAGES 7374–7383

DOI 10.1074/jbc.A109.058503

**Nickel ions inhibit histone demethylase JMJD1A and DNA repair enzyme ABH2 by replacing the ferrous iron in the catalytic centers.**

Haobin Chen, Nitai Charan Giri, Ronghe Zhang, Kenichi Yamane, Yi Zhang, Michael Maroney, and Max Costa

There were several errors in this article. The Coomassie blue stain of Flag-JMJD1A in [supplemental Fig. S2](#) was inadvertently duplicated in Fig. 1A. Additionally, the highest dose of  $\text{Ni}^{2+}$  used in Fig. 1A should read 250  $\mu\text{M}$  instead of 150  $\mu\text{M}$ . These errors have now been corrected and do not affect the results or conclusions of this work.




Fig. 1A

**Nickel ions inhibit histone demethylase JMJD1A and DNA repair enzyme ABH2  
by replacing the ferrous iron in the catalytic centers.**

Haobin Chen, Nitai Charan Giri, Ronghe Zhang, Kenichi Yamane, Yi Zhang, Michael Maroney and Max Costa

*J. Biol. Chem.* 2017, 292:10743.  
doi: 10.1074/jbc.A109.058503

---

Access the most updated version of this article at <http://www.jbc.org/content/292/25/10743>

Alerts:

- [When this article is cited](#)
- [When a correction for this article is posted](#)

[Click here](#) to choose from all of JBC's e-mail alerts

This article cites 0 references, 0 of which can be accessed free at  
<http://www.jbc.org/content/292/25/10743.full.html#ref-list-1>