

DELETION OF INTERLEUKIN 1 RECEPTOR ASSOCIATED KINASE 1 (*IRAK1*) IMPROVES GLUCOSE TOLERANCE PRIMARILY BY INCREASING INSULIN SENSITIVITY IN SKELETAL MUSCLE

Xiao Jian Sun^{##*}, Soohyun Park Kim[#], Dongming Zhang^{#&}, Helen Sun[#], Qi Cao[@], Xin Lu[@], Zhekang Ying[%], Liwu Li[¥], Robert R. Henry^{+¶}, Theodore P. Ciaraldi^{+¶}, Simeon I. Taylor[#], Michael J. Quon[#]

From the [#]Division of Endocrinology, Diabetes, and Nutrition, and [%]Division of Cardiovascular Medicine, Department of Medicine, and [@]Department of Diagnostic Radiology and Nuclear Medicine, University of Maryland School of Medicine, Baltimore, MD 21201, USA; ^{\$}Geriatric Research Education and Clinical Center, Baltimore Veterans Affairs Medical Center, Baltimore, MD 21201, USA; [&]The Second Affiliated Hospital of Zhengzhou University, Zhengzhou, China 450014; [¥]Virginia Tech, Blacksburg, VA 24061, USA; ⁺Veterans Affairs San Diego Healthcare System, San Diego, CA 92166, USA; [¶]Division of Endocrinology & Metabolism, University of California, San Diego School of Medicine, La Jolla, CA 92093, USA

Running Title: *Irak1* k/o mice have improved glucose tolerance

*To whom correspondence should be addressed: University of Maryland, Baltimore, Division of Endocrinology, Diabetes and Nutrition, 660 W. Redwood St., HH 594, Baltimore, MD 21201, USA. Tel: 410-706-5865; Fax: 410-706-1622; E-Mail: xsun@som.umaryland.edu.

Keywords: Inflammation, IRAK-1, insulin resistance, Akt, glucose tolerance

Chronic inflammation may contribute to insulin resistance (IR) via molecular cross-talk between pathways for pro-inflammatory and insulin signaling. Interleukin 1 receptor associated kinase 1 (IRAK1) mediates pro-inflammatory signaling via IL-1 receptor/Toll-like receptors, which may contribute to insulin resistance, but this hypothesis is untested. Here, we used male *Irak1* null (k/o) mice to investigate the metabolic role of IRAK1. C57BL/6 wild-type (WT) and k/o mice had comparable body weights on low-fat and high-fat diets (LFD and HFD, respectively). After 12 weeks on LFD (but not HFD), k/o mice (vs. WT) had substantially improved glucose tolerance [assessed by the intraperitoneal glucose tolerance test (IPGTT)]. As assessed with the hyperinsulinemic-euglycemic glucose clamp technique, insulin sensitivity was 30% higher in the *Irak1* k/o mice on chow diet, but the *Irak1* deletion did not affect IPGTT outcomes in mice on HFD, suggesting that the deletion did not overcome the impact of obesity on glucose tolerance. Moreover, insulin-stimulated glucose-disposal rates were higher in the k/o mice, but we detected no significant difference in hepatic glucose production rates (\pm insulin infusion). Positron emission/computed tomography scans indicated higher insulin-stimulated glucose uptake in muscle, but not liver, in *Irak1* k/o mice *in vivo*. Moreover, insulin-stimulated phosphorylation of Akt was higher in muscle, but not in liver, from *Irak1* k/o mice *ex vivo*. In conclusion, *Irak1* deletion improved muscle insulin sensitivity, with the effect being most apparent in LFD mice.

Insulin resistance is defined as decreased sensitivity or impaired responsiveness to metabolic actions of insulin

(1,2). In humans, insulin resistance is typically associated with impairments in multiple metabolic processes: insulin-stimulated glucose uptake in skeletal muscle and fat cells, antilipolysis in adipose tissue, and suppression of gluconeogenesis in liver (3-7). Studies have revealed many mechanisms that can induce insulin resistance. These include pro-inflammatory signaling pathways in cells (8-11), animals (12,13), and humans (14-16). Insulin regulates energy metabolism through a well-defined signaling pathway. As a consequence of binding to the insulin receptor (IR), insulin activates the receptor's intrinsic tyrosine kinase, which in turn catalyzes phosphorylation of intracellular substrates including insulin receptor substrate (IRS) proteins (17-19). Phosphotyrosine residues in IRS-proteins provide docking sites for SH2 domain-containing signaling molecules including PI3K (20-22) and Grb-2/mSOS, (20) leading to activation of PI3K and MAPK pathways. Lipid products of PI3K activate PDK-1, which triggers phosphorylation and activation of downstream serine kinases including Akt and PKC- ζ , enzymes with critical roles in mediating the metabolic actions of insulin (19,23). Proinflammatory cytokines including IL-1, and free fatty acids (FFA), activate innate immune signaling to NF- κ B via IL-1- and Toll-like receptors, which may impair insulin action in metabolic target tissues (24-26). This is mediated, in part, through serine phosphorylation of insulin receptor substrate-1 (IRS-1) by IKK β , JNK, and IRAK-1, which impairs insulin signaling, thereby inducing insulin resistance (11,16,27-33). Understanding the detailed mechanisms of this cross-talk between pro-inflammatory signaling pathways and insulin signaling pathways may help to improve the diagnosis of

prediabetic states, and also suggest new strategies to treat obesity and diabetes.

IRAK-1 is homologous to Pelle in *Drosophila* (34) and a key signaling molecule in mediating the inflammatory responses to IL-1-R and TLR. IL-1-R and Toll-like receptors interact with MyD88 to activate IRAK-4 which phosphorylates and activates IRAK-1 (26). Downstream from IRAK-1, TRAF6 interacts with TAK1 to activate the IKK complex leading to phosphorylation of I κ B α that then dissociates from NF- κ B (35,36). NF- κ B enters the nucleus to promote transcription of many pro-inflammatory genes including interferons and cytokines, such as IL-1 and TNF- α (37-39). IL-1, TLR-2/-4 receptor ligands (e.g., FFA) and TNF- α may cause insulin resistance in metabolic targets via IRS-1 serine phosphorylation catalyzed by IKK β , IRAK-1, and other innate immune kinases (28,40-44). Inhibiting IKK β reduces serine phosphorylation of IRS-1, which in turn ameliorates insulin resistance due to obesity or high fat diet (16,32,45-47). High activity genetic variants in the human *IRAK1* gene are associated with higher circulating C-reactive protein (a biomarker for heart disease risk and possibly diabetes) (48). IRAK-1 phosphorylates IRS-1 at Ser²⁴, which impairs binding of IRS-1 to PI3K, thereby impairing metabolic actions of insulin including insulin-stimulated translocation of GLUT4 (28). Interestingly, IRS-2 does not have a corresponding IRAK-1 phosphorylation site (27). Thus, IRAK-1 is not predicted to alter IRS-2 function. While IRAK-1 may impair metabolic actions of insulin primarily in cells (e.g., muscle) where IRS-1 is the predominant insulin signaling molecule downstream from the insulin receptor, the role of IRAK-1 might be less important in cells such as hepatocytes where IRS-2 plays the major role in mediating insulin actions (49-52).

In this study, we investigated the role of IRAK-1 in regulating glucose metabolism using *Irak1* null mice. We showed that *Irak1* null mice on low fat diet (but not on high fat diet that induces obesity, insulin resistance and glucose intolerance) have improved glucose tolerance likely mediated by increased insulin sensitivity in skeletal muscle but not in liver. In the context of precision medicine, these data suggest that low IRAK-1 activity in skeletal muscle may have predictive value to identify a subset of insulin resistant individuals most likely to obtain metabolic benefit from other therapeutic and lifestyle interventions that improve insulin sensitivity.

RESULTS

***Irak1* k/o mice display improved glucose tolerance on low fat, but not on high fat diet** - To investigate whether IRAK-1 deficiency alters glucose metabolism, we studied mice fed for 14 weeks with either low fat (LFD, 10% kcal from fat) or high fat diet (HFD, 60% kcal from fat). As expected, wild type mice fed with high fat diet had higher caloric intake (Fig. S1B and S1D) and also exhibited significantly higher body weight (Fig. S1A and S1C). While k/o mice ate slightly less food than WT mice

on either diet (4.6% less on LFD and 8.9% less on HFD), there were no significant differences in body weight between *Irak1* k/o and WT littermates on either diet (Fig. S1A and S1C). These observations suggest that *Irak1* k/o mice may have lower energy expenditure than WT littermates.

To investigate whether IRAK-1 deficiency altered glucose metabolism, we performed IPGTTs at baseline and also after 12 weeks of either HFD or LFD. There was no significant difference in baseline IPGTTs on normal chow diet between WT and k/o mice (data not shown). As expected, HFD induced glucose intolerance in both WT and k/o mice (Fig. 1C and 1D). Furthermore, there was no significant difference between WT and k/o mice with respect to IPGTTs on the HFD which induced obesity. By contrast, k/o mice had significantly enhanced glucose tolerance when fed a LFD (Fig. 1A and 1B). Therefore, although IRAK-1 deficiency improved glucose tolerance in mice fed with a LFD, it did not protect mice from HFD-induced obesity and impairment in glucose tolerance.

To investigate whether IRAK-1 deficiency affected insulin sensitivity, we performed insulin tolerance tests (ITT). Based on studies with tissue selective insulin receptor knockout mice, the ITT is primarily a reflection of insulin sensitivity in the liver rather than muscle (49,50). As expected, as judged by responses to ITT, both k/o and WT mice were less insulin sensitive when studied on HFD as compared to LFD (Figs. S2A - S2D). However, there were no significant differences between *Irak1* k/o mice and wild type littermates when compared with each other on either the HFD or the LFD (Figs. S2A - D). As demonstrated previously in studies with tissue-specific knockouts, deletion of insulin receptors in the liver (but not in muscle) impair the ability of insulin to lower plasma glucose levels (49,50). Thus, the ITT reflects primarily hepatic insulin sensitivity. Taken together, these data suggest that deletion of *Irak1* did not protect mice from HFD-induced hepatic insulin resistance characteristic of obesity.

***Irak1* k/o mouse displayed improved insulin sensitivity, and lowered markers of endothelial dysfunction and insulin resistance** - While ITTs reflect primarily hepatic insulin sensitivity, hyperinsulinemic euglycemic glucose clamp studies are the reference standard for *in vivo* measurement of insulin sensitivity in extrahepatic tissues including skeletal muscle and fat (1,53,54). Since we did not detect effects of *Irak1* deletion in mice fed a HFD, we conducted clamp studies in mice fed a normal chow diet. Average glucose was clamped at a steady-state level of 125 mg/dL by adjusting the glucose infusion rate (GIR) appropriately. *Irak1* k/o mice required significantly higher glucose infusion rates when compared with WT littermates to maintain steady-state blood glucose levels (Fig. 2A - 2B). After normalization for body weight (Fig. 2C) and steady-state blood glucose levels, we calculated an Insulin Sensitivity Index (ISI = GIR/steady-state glucose) (Fig. 2D). The ISI was higher for *Irak1* k/o mice than for their WT littermates when fed

with a normal chow diet. Thus, deletion of *Irak1* improved insulin sensitivity in extrahepatic tissues under these conditions (Fig. 2).

IRAK-1 is a key signaling molecule in TLR/IL1-R/TNF α -R mediated inflammatory responses and may contribute to the pathogenesis of chronic inflammation-associated insulin resistance. To explore this issue, we measured insulin resistance, cardiometabolic and inflammation biomarkers in mice fed for 14 weeks with a LFD and HFD. Fasting (5 h) plasma glucose was significantly lowered in *Irak1* k/o mice as compared with wild type mice fed with LFD (Fig. 3B). Although fasting plasma insulin concentration tended to be lower in *Irak1* k/o mice, this did not reach statistical significance (Fig. 3A). Nevertheless, QUICKI, a fasting surrogate index for insulin sensitivity (55,56), was significantly higher (Fig. 3C), suggesting an improvement of insulin sensitivity for k/o mice when compared with wild type littermates. There was no difference in adiponectin levels between k/o and wild type littermates (Fig. 3D). ET-1 is a potent vasoconstrictor and a biomarker for endothelial dysfunction associated with insulin resistance (59-61). Interestingly, deletion of *Irak1* led to lowered ET-1 levels (Fig. 3E) suggesting an improvement in endothelial function in *Irak1* k/o mice consistent with a role for inflammation in endothelial dysfunction related to insulin resistance. TNF- α is a biomarker for inflammation, and is reported to be associated with insulin resistance (57,58). Plasma levels of TNF- α in our experiments were too low to obtain reliable data with our immunoassay although there was a trend toward decreased levels of TNF- α in k/o mice as compared to WT mice (data not shown), consistent with attenuation of chronic inflammation.

As expected, when compared with low fat diet, high fat diet mice showed higher insulin resistance markers including fasting insulin (Fig. 3G), glucose (Fig. 3H), ET-1 (Fig. 3K), and lower adiponectin (Fig. 3J) and QUICKI (Fig. 3I). This confirmed that 14 weeks of high fat diet induced obesity (Fig S1, C and D) and insulin resistance. However, we did not observe a significant difference in ET-1 (Fig. 3K) between WT and k/o as was seen in the low fat diet study (Fig. 3E). Adiponectin (Fig. 3J) and CRP (Fig. 3L) also showed no difference between WT and k/o mice. Of note, we observed lower plasma insulin levels in k/o mice (Fig. 3G), and, as a result, higher QUICKI (Fig. 3I). These data are consistent with the conclusion that the k/o were more insulin sensitive than WT mice.

Improved insulin sensitivity in *Irak1* k/o mice may be caused specifically by increased glucose uptake in skeletal muscle - To assess contributions of specific tissues to improvement in glucose metabolism *in vivo*, we conducted ^{18}F -FDG PET/CT scans to investigate insulin-stimulated ^{18}F -FDG bio-distribution in three insulin sensitive target tissues: hind limb muscle, abdominal subcutaneous white adipose tissue, and liver (Fig.S3). We did not observe higher ^{18}F -FDG uptake in the liver post insulin injection in either *Irak1* k/o or wild type mice (Fig.

4A - 4B). This suggests that altered liver glucose metabolism did not account for improved insulin sensitivity observed in *Irak1* k/o mice. By contrast, in skeletal muscle, ^{18}F -FDG uptake was significantly higher in k/o mice when compared with wild type littermates. This achieved statistical significance at 20-30 min (i.e., the delayed wash-out phase of ^{18}F -FDG), but not in the vascular phase (i.e., the 0-10 min time frame) (Fig. 4A - 4B). These results demonstrated an important role of IRAK-1 in skeletal muscle glucose uptake, and likely underlies the increased insulin sensitivity observed in *Irak1* k/o mice on LFD. White adipose tissue of the abdomen also showed a trend toward delayed wash-out of ^{18}F -FDG after insulin injection in both types of animals, but there was no difference between k/o and wild type mice (Fig.4A - B). Thus, it is unlikely that adipose tissue had an important direct role in mediating the beneficial effect of IRAK-1 deficiency to improve insulin sensitivity.

To further support these findings, we performed hyperinsulinemic euglycemic glucose clamp studies with [3-H^3]-glucose tracer on *Irak1* k/o and WT littermates fed with chow diets. The hyperinsulinemic euglycemic clamp with [3-H^3]-glucose tracer is the reference standard method for assessing insulin sensitivity in muscle, fat, and liver (1,53,62). As expected, *Irak1* k/o mice required significantly higher glucose infusion rate resulting in a larger insulin sensitivity index (ISI) as compared to wild type littermates (Fig. 5A and 5B). Glucose disposal rates were comparable at the basal state, but significantly higher in *Irak1* k/o mice during insulin infusion (Fig. 5C). When hepatic glucose production was measured, there was no significant difference between *Irak1* k/o mice and their WT littermates in either the basal state or the insulin-stimulated state (Fig. 5D). These data are consistent with the PET/CT scan data showing improved insulin sensitivity in skeletal muscle but not liver. This suggests that improvement of insulin sensitivity in *Irak1* k/o mice mainly occurs in muscle and/or adipose tissue when compared with wild type mice.

IRAK-1 deficiency is associated with improved intracellular insulin signaling in skeletal muscle and adipose tissue, but not in liver - Cross-talk between the pathways of inflammatory signaling and insulin signaling is likely mediated, at least in part, by serine/threonine phosphorylation of IRS-1, which attenuates insulin signaling, thereby inducing insulin resistance (27). Based on this hypothesis, IRAK-1 deficiency is predicted to alleviate the negative impact of inflammation upon insulin signaling and action. To evaluate this issue, we investigated the effect of IRAK-1 deficiency on the insulin signaling pathways in metabolic insulin target tissues including skeletal muscle, adipose tissue, and liver. First, we measured and compared IRAK-1 protein levels in these tissues from wild type mice (Fig. S4A). Quantified by total proteins of tissue lysates, muscle expressed higher IRAK-1 protein (100%) than liver (31%) or fat (42%). Second, we measured IRAK-1 expression in

these tissues of WT and k/o mice. As expected, IRAK-1 was markedly lower in all the tissues tested from *Irak1* k/o mice (Fig. S4B). There was a weak band in the *Irak1* k/o muscle sample (Fig. S4B, top panel), but not in fat tissue or liver (Fig. S4B, middle and lower panels). Although the observed weak band in muscle immunoblots are likely non-specific, we cannot entirely rule out the possibility that there is less than complete knockout of IRAK-1 in skeletal muscle.

As expected, insulin stimulation showed higher phosphorylation of Akt in all three tissues. Phosphorylation is a proxy for activation of the enzyme by insulin in skeletal muscle, liver, and adipose tissue (Fig. 6). Interestingly, IRAK-1 deficient skeletal muscle showed significantly higher insulin-stimulated phosphorylation of Akt when compared with that of wild type littermates (Fig. 6A – 6B). Whereas fat tissue isolated from *Irak1* k/o mice showed a trend toward higher insulin stimulated Akt kinase activity, this did not reach statistical significance (Fig. 6E – 6F). By contrast, liver from *Irak1* k/o mice did not show any difference in insulin stimulated Akt phosphorylation when compared with wild type littermates (Fig. 6C – 6D). This is consistent with *in vivo* data from both glucose clamp and PET/CT studies showing comparable liver insulin sensitivity in *Irak1* k/o and WT mice.

DISCUSSION

Protein phosphorylation plays multiple roles related to insulin action and insulin resistance. The insulin receptor possesses intrinsic tyrosine kinase activity, which catalyzes tyrosine phosphorylation of IRS-1, IRS-2, and other endogenous substrates. This in turn triggers signaling pathways that activate downstream serine-specific protein kinases that mediate insulin action (e.g., PDK-1 and Akt). On the other hand, some serine-specific protein kinases participate in multiple biological pathways and are able to serine phosphorylate IRS-1 and IRS-2. For example, some inflammatory signaling pathway kinases (e.g., IKK, JNK, MAPK, and IRAK-1) have been reported to attenuate insulin signaling and induce insulin resistance (27) by promoting serine-specific phosphorylation of components of the insulin signaling pathway (e.g., IRS-1 and/or IRS-2). This has led to the hypothesis that chronic inflammation represents a key mechanism contributing to the pathogenesis of insulin resistance in obesity (63,64).

Our current studies suggest that suppression of chronic inflammation may be beneficial by ameliorating insulin resistance. IRAK-1 is a critical enzyme in signaling pathways downstream of the IL-1 receptor TNF- α receptor, and TLRs. In the present study, we demonstrated that deletion of *Irak1* improved insulin sensitivity in mice fed a LFD (but not with HFD-induced obesity). Our observations suggest that the improvement in whole body insulin sensitivity is most likely mediated by improved insulin signaling and enhanced insulin sensitivity in skeletal muscle. The significance of this study is that although IRAK-1 is missing in all tissues, it appears that

muscle may be the tissue in which IRAK-1 exerts the most important metabolic effects. There may also be secondary metabolic benefits mediated by IRAK-1 absence in white fat, and vascular endothelium (28,65).

Role of IRAK-1 in inflammation-associated insulin resistance - IRAK-1 deficient mice, which have been studied for over a decade, are characterized by a partial impairment in inflammatory response to LPS at low dose but normal response to high dose of LPS (66,67). Despite complete absence of IRAK-1, complete functional impairment has only been observed for a limited number of responses – e.g., interferon- α production (68). Although adverse metabolic effect of LPS in IRAK-1 k/o mice have been studied (69), the metabolic phenotype of *Irak1* k/o mice in response to insulin has not been reported previously. Thus, inspired by our previous observation that IRAK-1 mediated phosphorylation exerts a negative impact on IRS1-mediated insulin signaling (28), the present set of experiments was designed to investigate the role of IRAK-1 in metabolic physiology. Specifically, what is the role of IRAK-1 in the pathogenesis of the insulin resistant state associated with chronic inflammation? Importantly, deletion of *Irak1* did not affect body weight – regardless of the fat content of the diet (i.e., LFD vs. HFD). Because body weight was not affected by deletion of *Irak1*, this eliminated the confounding influence of obesity. Interestingly, deletion of *Irak1* was associated with a “healthy phenotype” (i.e., supra-normal insulin sensitivity) in mice fed with LFD. This was an unexpected finding as it suggests that IRAK-1 appears to be important under conditions associated with a minimal level of inflammation. By contrast, both genotypes exhibited comparable degrees of insulin resistance when fed a HFD. Taken together, these data suggest that diet-induced obesity essentially eliminated the supra-healthy phenotype associated with *Irak1* k/o genotype. This is most likely because there are multiple mechanisms for obesity to cause insulin resistance. Therefore, in the presence of obesity-induced insulin resistance the contribution of IRAK-1 alone is masked.

High fat diets have been widely used to induce obesity in animal models as well as to study mechanisms of insulin resistance (70). Multiple factors have been reported to contribute to the molecular pathogenesis of obesity-induced insulin resistance. In addition to chronic inflammation (11,71), other pathogenic factors including hyperlipidemia (72,73), ectopic lipid accumulation (74), gut hormones (75), oxidative stress (76), ER stress (9,77), sympathetic nervous system dysfunction (78), and gut microbiota (79) also contribute to insulin resistance in obesity. In this context, it is not surprising that *Irak1* deficiency did not protect mice from obesity-induced insulin resistance inasmuch as IRAK-1 is only one of many mechanisms contributing to induction of insulin resistance. Low fat and normal chow diets are likely associated with minimal “low grade” inflammation. Nevertheless, when this low grade inflammation was suppressed by deleting *Irak1*, this enhanced insulin

sensitivity and glucose tolerance significantly. Our observations in metabolic physiology mirror what has been reported in the context of the physiology of inflammation. Specifically, macrophages from *Irak1* k/o mice have impaired production of TNF- α in response to low concentrations of LPS (a ligand for TLR4 and the TNF- α receptor upstream of IRAK-1). However, this defect can be overcome by high concentrations of LPS, which induce the same quantities of TNF- α in macrophages regardless of *Irak1* genotype. Thus, *Irak1* deletion only partially impairs the inflammation pathway induced by IL-1/TLR signaling (66,67). Both IL-1 and TNF- α stimulate increased phosphorylation of IRS-1 (Ser²⁴) in isolated rat adipose cells (28). These previously published data provide a precedent to explain why *Irak1* k/o mice exhibit improved insulin sensitivity when fed either a LFD or normal chow diet (i.e., low level inflammation) while the protective effect is overwhelmed on a HFD (i.e., high level of inflammation). Interestingly, although k/o mice have slightly lower food intake (4.5% on LFD and 8.9% on HFD) than that of WT mice, they maintained the same body weight as WT mice. This suggests lower metabolic rate in k/o mice counterbalances the lower food intake. Further study will clarify whether the apparent decrease in energy utilization is mediated by the central nervous system or peripheral tissues.

IRAK-1 is not the only kinase that phosphorylates IRS-1 (Ser²⁴). We and others reported that a number of PKC isoforms also phosphorylate IRS-1 (Ser²⁴) to impair metabolic insulin signaling (80,81). Therefore, it is possible that IRS-1 Ser²⁴ may be phosphorylated by other kinases in mice where IRAK-1 is missing. However, the fact that we observed a significant and substantial metabolic phenotype in *Irak1* k/o mice consistent with abrogation of cross-talk between innate immune signaling and insulin signaling suggests that the contribution of IRAK-1 to regulate whole body glucose homeostasis is important relative to other kinases that also phosphorylate IRS-1 (Ser²⁴).

Interestingly, fasting levels of plasma ET-1, a potent vasoconstrictor, were also significantly lower in k/o mice vs. WT ($p = 0.016$) after 14 weeks of LFD (20 week old mice). Reduction of plasma ET-1 may be secondary to the reduction of TNF- α which is a positive regulator of ET-1 expression. ET-1 plays an important role in the pathogenesis of insulin resistance-associated endothelial dysfunction (24,60,82). However, there is also likely a direct effect of IRAK-1 absence in endothelial cells to reduce ET-1 synthesis and secretion. In a previous study, we found that siRNA knockdown of TLR2 (closely upstream of IRAK-1) in primary vascular endothelial cells completely abrogates the effect of FFA (TLR2 ligand elevated in insulin resistance) to stimulate phosphorylation of IRS-1 (Ser²⁴) (65). Moreover, under conditions of metabolic insulin resistance (i.e., pathway-selective impairment of PI 3-kinase-dependent insulin signaling with intact MAPK signaling), the effect of insulin to stimulate increased synthesis and secretion of

ET-1 from vascular endothelial cells is greatly enhanced (83-85). Taken together, these data suggest that the ability of IRAK-1 to phosphorylate IRS-1 (Ser²⁴) in endothelium may promote vascular insulin resistance leading to endothelial dysfunction characterized by increased insulin- and FFA-stimulated ET-1 synthesis and secretion. This is relevant because whole body metabolic insulin resistance is characterized by elevated fasting and post-prandial insulin and FFA levels as well as pathway-selective impairment in PI 3-kinase-dependent insulin signaling and action. We hypothesize that decreased levels of ET-1 mediated directly and indirectly by absence of IRAK-1 may improve endothelial function and increase blood flow to skeletal muscle. This will increase access of insulin and glucose to metabolic tissues and secondarily promote insulin-stimulated glucose uptake and improved glucose tolerance (61).

Role of skeletal muscle in mediating physiological effects of IRAK-1 deficiency - Several lines of evidence suggest that deletion of IRAK-1 increased insulin sensitivity and improved glucose metabolism in skeletal muscle, but not in liver.

1. Hyperinsulinemic euglycemic glucose clamp studies demonstrated that IRAK-1 deficiency was associated with higher insulin-stimulated glucose disposal rate, but did not alter insulin's ability to suppress the rate of hepatic glucose production. Insulin-stimulated glucose disposal as assessed by clamp studies has been demonstrated to reflect primarily glucose uptake by skeletal muscle. Observations in both mice and humans suggest that insulin tolerance tests (ITT) primarily reflect hepatic insulin sensitivity. Whereas muscle-specific insulin receptor knockout mouse exhibit normal ITT (49), insulin tolerance was impaired in mice with liver-specific insulin receptor knockout mouse (50). Similarly, in patients with insulinoma, insulin-driven fasting hypoglycemia is mediated primarily by suppression of hepatic glucose production (86). Thus, the normal ITT observed in *Irak1* k/o mice support the conclusion that the liver responds normally to insulin.
2. In PET/CT studies, muscle and adipose tissue were the principal tissues into which insulin promoted distribution of ¹⁸F-FDG. Further, deletion of IRAK-1 selectively increased muscle ¹⁸F-FDG uptake. There was no difference in liver ¹⁸F-FDG uptake.
3. Deletion of *Irak1* enhanced insulin signaling in muscle, but not liver, as assessed by analysis of insulin-stimulated phosphorylation of Akt (Fig. 6) and MAP kinase (Data not shown). There was also a trend toward improved insulin signaling in adipose tissue of k/o mice (although data did not achieve statistical significance).

The PET/CT scan does not provide a direct assessment of glucose uptake, but rather relies on of radioactively-labeled 2-deoxy-2-fluoro-D-glucose molecule (FDG) uptake as tracer. Because fasting glucose was lower in k/o mice than controls, this could differentially affect 2-deoxy-2-fluoro-D-glucose molecule (FDG) tracer uptake in different tissues. As compared to glucose, the 2-

hydroxyl group in FDG is replaced by a fluorine atom. While FDG does undergo phosphorylation on the 6-hydroxyl, it is not a substrate for downstream steps in the glycolytic pathway. Accordingly, the rate of accumulation of FDG is determined by the rate of glucose transport, which is mediated primarily by GLUT-family transporters. FDG uptake by the liver is mediated primarily by GLUT2, which has a K_m for glucose of ~17 mM (101). As shown in Fig. 3B, fasting glucose levels (~10 mM) are below the K_m of GLUT2 to transport glucose, and the majority of GLUT2 molecules would be available to transport FDG with relatively little competition from unlabeled glucose. Accordingly, FDG uptake by the liver would be predicted to be relatively insensitive to ambient glucose levels (for glucose levels substantially below the K_m of GLUT2 for glucose). In this context, it is noteworthy that both WT and k/o mice exhibit similar rates of FDG uptake in liver (Fig. 4).

By contrast, the K_m for GLUT1 is ~26 mM and for GLUT4 is ~4-5 mM (87). Thus, for ambient glucose levels of ~10 mM, approximately two thirds of GLUT4 molecules would be engaged in transporting glucose while the majority of GLUT1 molecules would not be bound to glucose molecules. While GLUT4 is believed to be the principal glucose transporter in both adipose tissue and skeletal muscle in the presence of insulin, both GLUT1 and GLUT4 are believed to be significant in the absence of insulin. While GLUT1-mediated FDG transport would be predicted to be relatively insensitive to inhibition by glucose levels substantially below the K_m =26 mM, GLUT4-mediated FDG transport would likely be significantly inhibited by glucose levels of ~10 mM. Thus, it is possible that GLUT1 would be the principal transporter mediating FDG uptake when plasma glucose levels are ~10 mM in the absence of insulin. In any case, ambient glucose levels would be predicted to exert similar effects on FDG uptake in both adipose tissue and skeletal muscle. In this context, it is noteworthy that rates of FDG uptake in adipose tissue are not significantly different in WT vs. IRAK-1 knockout mice (Fig. 4). In contrast, FDG uptake by skeletal muscle is selective increased in knockout mice. If the difference observed for skeletal muscle were caused primarily by competitive inhibition at the level of GLUT4, one might have expected to observe a similar difference for adipose tissue, which is not the case.

Taken together, these *in vivo* and *ex vivo* analyses suggest that IRAK-1 plays a particularly important role in the metabolic physiology of skeletal muscle, but little, if any, effect in liver. What are the mechanisms whereby deletion of *Irak1* selectively promotes insulin action in muscle rather than liver? Several possibilities deserve consideration. At least in theory, this might be explained by tissue selective expression of IRAK-1. Although our western blot analyses for muscle, fat and liver tissues demonstrate that all three tissues express IRAK-1, relative expression of IRAK-1 is higher in muscle than fat and liver. A second possible explanation is based on tissue-

specific roles of insulin receptor substrate molecules. Whereas muscle and adipocytes rely primarily on IRS-1, IRS-2 is more important in liver. As reported previously, IRAK-1 phosphorylates Ser²⁴ on IRS-1. Phosphorylation of Ser²⁴ impairs the signaling function of IRS-1 (28). By contrast, the Ser²⁴ phosphorylation site is not conserved in the IRS-2 molecule (27). Thus, deletion of *Irak1* has potential to improve insulin sensitivity in tissues where IRS-1 is important (e.g., skeletal muscle), but is less likely to affect insulin signaling in tissues such as liver where IRS-2 has the dominant role in insulin signaling. Indeed, consistent with this possibility, our observations from clamp studies demonstrated that insulin-stimulated inhibition of hepatic glucose production was comparable among *Irak1* k/o and WT mice on LFD while insulin sensitivity in skeletal muscle was improved in the *Irak1* k/o mice when compared with WT mice. Furthermore, there is a precedent for other kinases regulating insulin signaling to phosphorylate IRS proteins in an isoform specific manner due to the presence of a phosphorylation site in one isoform but not another. For example, PKC- ζ phosphorylates IRS-1 but not IRS-2 (88). This IRS isoform-specific phosphorylation may represent a mechanism for determining insulin signaling specificity in different tissues. In immunoblotting studies, we did not detect the presence of phospho-IRS-1 (Ser²⁴) in skeletal muscle in either the absence or presence of insulin stimulation under conditions of our *ex vivo* insulin signaling studies (in either in wild type or k/o mice) (data not shown). Although we cannot entirely eliminate the possibility of technical problems, we used a phospho-specific antibody that we previously developed and characterized that detected phospho-IRS-1 (Ser²⁴) in rat fat cells (28) and bovine aortic endothelial cells (65) *in vitro*. Taken at face value, these data suggest that Ser²⁴ phosphorylation of IRS-1 may not be important in mouse skeletal muscle. Nevertheless, it is also possible that phosphorylation of Ser²⁴ might accelerate degradation of IRS-1, thereby decreasing the level of phospho-IRS-1 (Ser²⁴) to the point where it is not detectable by immunoblotting. On the other hand, our immunoblotting experiments did not detect an alteration in the levels of IRS1 in IRAK1-knockout mice (data not shown). Another possibility is that rapid dephosphorylation of phospho-IRS-1 (Ser²⁴) is occurring in our *ex vivo* experimental setting such that we were unable to detect a transient but physiologically relevant higher phospho-IRS-1 (Ser²⁴) in skeletal muscle by immunoblotting.

A third possibility to explain why deletion of *Irak1* selectively promotes insulin action in muscle rather than liver relates to the cell biology of insulin action in different tissues. In muscle cells, insulin-stimulated glucose uptake is mediated by insulin-stimulated translocation of GLUT4-containing vesicles from an intracellular location to the plasma membrane (89,90). By contrast, GLUT2-mediated transport of glucose into hepatocytes is not directly regulated by insulin. Moreover, deletion of *Irak1* did not alter levels of GLUT4 protein in muscle (Fig.

S4C). Thus, we hypothesize that *Irak1* deletion leads to diminished serine phosphorylation of IRS-1, which in turn promotes the activity of the GLUT4 translocation machinery, thereby increasing insulin sensitivity (28,89). However, as noted above, insulin-stimulated hepatic glucose production was comparable between *Irak1* k/o mice and WT mice making it more likely that it is the predominance of IRS-2 in liver and the lack of an IRAK phosphorylation site on IRS-2 that explains why liver insulin sensitivity did not change in *Irak1* k/o mice. Thus, the improvement in glucose tolerance observed in *Irak1* k/o mice on LFD is likely mediated primarily by improved metabolic insulin signaling and action in muscle.

Conclusion: possible translational implications of role of IRAK-1 in metabolic physiology. Our studies in a rodent animal model suggest that IRAK-1 deficiency is associated with a healthy phenotype (i.e., supranormal insulin sensitivity) – at least in mice fed with either LFD or normal chow diet. Inasmuch as the *Irak1* genotype does not affect the degree of insulin resistance in mice fed a HFD, we hypothesize that *Irak1* k/o mice will experience a quantitatively greater deterioration of insulin sensitivity when switched from normal chow to a HFD. Conversely, *Irak1* k/o mice might experience greater metabolic benefit (i.e., a quantitatively greater improvement in insulin sensitivity) if weight loss were induced by shifting the mice from a HFD to a LFD. If our data in this animal model are predictive of human pathophysiology, this prompts the hypothesis that IRAK-1 activity in skeletal muscle may provide a biomarker to predict a beneficial metabolic response to weight loss. If correct, this hypothesis may have significant implications for a precision medicine approach to weight loss. In other words, low levels of skeletal muscle IRAK-1 activity might identify a subset insulin resistant patients who are most likely to experience the greatest improvement in insulin sensitivity in response to weight loss.

EXPERIMENTAL PROCEDURES

Animal studies: All animal protocols were approved by the Institutional Animal Care and Use Committee of the University of Maryland School of Medicine. To generate *Irak1* null mice and their wild type control, female homozygotes were first backcrossed to C57BL/6 males for 10 generations. Since the *Irak1* null allele was introduced on a genetic background of C57BL/6 mice (Harlan Laboratories), we genotyped for a previously reported mutation in nicotinamide nucleotide transhydrogenase (*Nnt*), and confirmed that our mice did not carry a mutation in *Nnt* (91,92). The *Irak1* gene is on the X-chromosome (93). Thus, males that are hemizygous for the null allele of *Irak1* will be deficient in *Irak1* function (k/o) whereas male's hemizygous for the functional allele of *Irak1* will be wild type (WT). To simplify breeding and selection of affected mice, we limited our studies to male mice (either *Irak1* k/o or *Irak1* WT). For studies on different diets, *Irak1* k/o vs. WT littermate mice were fed with LFD (10% kcal from fat,

70% kcal from Carbohydrate, #D12450B, Research Diets) or HFD (60% kcal from fat, 20% kcal from Carbohydrate, #D12492, Research Diets) for indicated weeks or chow diet (18% kcal from fat and 58% kcal from carbohydrate, #2018SX, Teklad Global). Body weight and food intake were monitored 3 times per week.

IPGTT and ITT: After 5 h fasting, either glucose (1 g/kg for IPGTT) or human insulin (0.75 IU/kg for ITT, Eli Lilly, Indianapolis, IN) was injected intraperitoneally (50). Blood glucose was monitored at indicated time points with an Accu-Chek Aviva Blood glucometer. Area under the curve (AUC) for IPGTT or area above the curve (AAC) for ITT were calculated based on the trapezoidal rule.

Hyperinsulinemic euglycemic glucose clamp studies: (53,94) Jugular veins were cannulated 4-5 day before the clamp experiments. Mice were acclimated to restrainers three times (3 h each time) before clamp. After 5 h fasting, [^3H]-glucose was infused intravenously (2.5 μCi bolus followed by constant infusion at 0.05 $\mu\text{Ci}/\text{min}$). Blood samples (20 μL) were collected at 90, 100, and 110 min after the initiation of ^3H -glucose infusion for the measurement of ^3H -glucose specific activity, blood glucose concentration, and calculation of basal hepatic glucose production. Then, regular human insulin was infused (60 mU/kg loading bolus followed by constant infusion at 2.5 mU/kg/min), and glucose levels were measured every 10 min. An infusion of 25% dextrose was adjusted to maintain the blood glucose at 120–130 mg/dL. Blood samples were again collected at 100, 110 and 120 min during insulin infusion for calculation of insulin stimulated glucose disposal rate and hepatic glucose production rate. Glucose infusion rate (GIR) was calculated as the average glucose infusion rate during the last 30 min of the clamp steady-state period and normalized for body weight. An insulin sensitivity index (ISI) was defined as GIR divided by the level at which glucose was clamped (GIR/clamp glucose). The rates of glucose disposal and hepatic glucose production were calculated as previously described (94).

Portal vein insulin injection and collection of liver, muscle, and fat tissues: Mice were fasted for 5 h and anesthetized with isoflurane vapor. After making a ventral incision to expose the liver and portal vein, we injected either human insulin (10 U/kg body weight, Eli Lilly, Indianapolis, IN) or saline into the portal vein (95-97). After 5 min, samples of epididymal fat tissue, liver, and hind limb skeletal muscle were collected and flash frozen for future immunoblotting analysis (95-97).

Tissue and cell sample preparation and immunoblotting: Muscle (200 mg) and liver (100 mg), frozen tissues were homogenized at 4 °C in 400 μL lysis buffer (50 mM HEPES, pH 7.4, 2 mM EDTA, 50 mM NaF, 100 mM NaCl, 5 mM DTT, 0.1% NP40, 2 mM Benzamidine, 0.1 mM Na_3VO_4 , and 5 μL of Roche protein inhibitor cocktail set) with a tissue grinder. Tissue lysates were centrifuged at 21,000 x g for 1 h at 4 °C in a table top centrifuge. The supernatants were used for

immunoblotting analysis. For fat tissue, 400 mg of frozen tissue was homogenized in 400 μ L of fat tissue lysis buffer (10 mM Tris, pH 7.4, 1 mM EDTA, 100 mM NaF, 0.2 mM Na_3VO_4 , and 4 μ L of Roche protease inhibitor cocktail set). Homogenates were mixed with 900 μ L of cold acetone at -80 °C overnight, and then centrifuged at 21,000 x g for 1 h at 4 °C. Supernatants were discarded and the pellets were dried on ice for 5-10 min prior to resuspension in 400 μ L of Laemmli sample buffer for immunoblotting analysis. Aliquots (20 μ g of total protein) from each tissue sample were separated on a 10% SDS PAGE gel, transferred to PVDF membranes, and incubated with indicated antibodies as previously described (95-97). Antibodies against IRAK-1, pAkt (S473), Akt, GLUT4, GAPDH and β -actin were purchased from Cell Signaling Technology Inc. Quantification of images was performed using an imaging device with Image Lab software. Relative pAkt was calculated by dividing pAkt density with that of total Akt. Relative IRAK-1 in muscle, liver and fat tissues were calculated by dividing all three tissues by muscle IRAK-1 density.

PET/CT scan: 3 *Irak1* k/o and 3 WT male mice were fasted overnight, and [^{18}F]-fluorodeoxyglucose (FDG) positron emission tomography-computed tomography (PET/CT) was performed with animals maintained under 2% isoflurane anesthesia on a Siemens Inveon Small Animal PET/CT Imaging System (Siemens Medical Solutions USA, Inc.) in the Core for Translational Research in Imaging (CTRIM). [^{18}F]-FDG (11.1 MBq; 300 μ Ci) and insulin (1.125 U/kg body weight) in 200 μ L of saline were injected via tail vein of anesthetized mice. Dynamic imaging was acquired for 30 min. Data acquisition, image reconstruction, and region of interest (ROI) analysis were carried out according to previously reported methods (98,99). The data were obtained by ROI analysis focused on several target organs: hind limb muscle, subcutaneous white adipose tissue (abdominal region), liver, and brain. The Quantitative [^{18}F]-FDG PET data in the target regions were expressed as percentages of injected dose per gram (%ID / g) at three time points of 0-10, 10-20 and 20-30 min.

ELISA: Adiponectin (Millipore), Endothelin-1 (Enzo

Life Science), insulin (Concodian), TNF- α (Invitrogen) and CRP (ThermoFisher Scientific) were measured by ELISA according to the manufacturers' instructions.

Statistical analyses: T-tests were performed with Excel software, and ANOVA for repeated measures was conducted with Axum 7 software.

Acknowledgements

This work was supported by grants from the American Diabetes Association to Dr. Quon (1-13-BS-150) and Dr. Sun (1-14-BS-198) as well as a grant from the National Institute of Environmental Health Sciences to Dr. Ying (R01ES024516). This work was supported in part by Merit Review Award (I01CX00635) from the United States Department of Veterans Affairs Clinical Sciences Research and Development Service. The contents do not represent the views of the U.S. Department of Veterans Affairs or the United States Government. This work was also supported by Baltimore Veterans Affairs Medical Center Geriatric Research Clinical Center (GRECC). In addition, Dr. Taylor acknowledges support from the National Institute of Diabetes and Digestive and Kidney Diseases (5P30DK072488-12).

Conflicts of Interest

SIT has worked as a paid consultant for Ionis Pharmaceuticals and Aegerion Pharmaceuticals. Other authors have nothing to disclose.

Author Contributions

XJS designed, performed and analyzed the experiments and coordinated the study and wrote the paper, SPK performed the animal care, IPGTT and ITT, ELISA measurement and immunoblotting analysis, DZ and HS participated in the immunoblotting of animal tissues, QC and XL performed and analyzed the experiments shown in Figure S3, ZY contributed to the discussion, LL contributed to the *Irak1* k/o mice, RRRH and TPC contributed to the interpretation of data and writing of the manuscript. ST contributed to the discussion of the project and writing the manuscript. MJQ conceived and coordinated the study, analyzed all data, wrote, and edited the manuscript.

REFERENCES

1. Muniyappa, R., Lee, S., Chen, H., and Quon, M. J. (2008) Current approaches for assessing insulin sensitivity and resistance in vivo: advantages, limitations, and appropriate usage. *Am J Physiol Endocrinol Metab* **294**, E15-26
2. Kahn, C. R. (1994) Banting Lecture. Insulin action, diabetogenes, and the cause of type II diabetes. *Diabetes* **43**, 1066-1084
3. Samuel, V. T., and Shulman, G. I. (2016) The pathogenesis of insulin resistance: integrating signaling pathways and substrate flux. *J Clin Invest* **126**, 12-22
4. Kim, Y. B., Kotani, K., Ciaraldi, T. P., Henry, R. R., and Kahn, B. B. (2003) Insulin-stimulated protein kinase C lambda/zeta activity is reduced in skeletal muscle of humans with obesity and type 2 diabetes: reversal with weight reduction. *Diabetes* **52**, 1935-1942
5. Matsumoto, M., Pocai, A., Rossetti, L., Depinho, R. A., and Accili, D. (2007) Impaired regulation of hepatic glucose production in mice lacking the forkhead transcription factor Foxo1 in liver. *Cell Metab* **6**, 208-216
6. Coppack, S. W., Jensen, M. D., and Miles, J. M. (1994) In vivo regulation of lipolysis in humans. *J Lipid Res* **35**, 177-193
7. Wahren, J., and Ekberg, K. (2007) Splanchnic regulation of glucose production. *Annu Rev Nutr* **27**, 329-345
8. Jaeschke, A., and Davis, R. J. (2007) Metabolic stress signaling mediated by mixed-lineage kinases. *Mol Cell* **27**, 498-508
9. Hotamisligil, G. S. (2010) Endoplasmic reticulum stress and atherosclerosis. *Nat Med* **16**, 396-399
10. Wellen, K. E., Fucho, R., Gregor, M. F., Furuhashi, M., Morgan, C., Lindstad, T., Vaillancourt, E., Gorgun, C. Z., Saatcioglu, F., and Hotamisligil, G. S. (2007) Coordinated regulation of nutrient and inflammatory responses by STAMP2 is essential for metabolic homeostasis. *Cell* **129**, 537-548
11. Hotamisligil, G. S. (2003) Inflammatory pathways and insulin action. *Int J Obes Relat Metab Disord* **27 Suppl 3**, S53-55
12. Fujimoto, Y., Torres, T. P., Donahue, E. P., and Shiota, M. (2006) Glucose toxicity is responsible for the development of impaired regulation of endogenous glucose production and hepatic glucokinase in Zucker diabetic fatty rats. *Diabetes* **55**, 2479-2490
13. Kewalramani, G., Bilan, P. J., and Klip, A. (2010) Muscle insulin resistance: assault by lipids, cytokines and local macrophages. *Curr Opin Clin Nutr Metab Care* **13**, 382-390
14. DeFronzo, R. A. (2010) Insulin resistance, lipotoxicity, type 2 diabetes and atherosclerosis: the missing links. The Claude Bernard Lecture 2009. *Diabetologia* **53**, 1270-1287
15. Hommelberg, P. P., Langen, R. C., Schols, A. M., Mensink, R. P., and Plat, J. (2010) Inflammatory signaling in skeletal muscle insulin resistance: green signal for nutritional intervention? *Curr Opin Clin Nutr Metab Care* **13**, 647-655
16. Shoelson, S. E., Lee, J., and Goldfine, A. B. (2006) Inflammation and insulin resistance. *J Clin Invest* **116**, 1793-1801
17. Sun, X. J., Rothenberg, P., Kahn, C. R., Backer, J. M., Araki, E., Wilden, P. A., Cahill, D. A., Goldstein, B. J., and White, M. F. (1991) Structure of the insulin receptor substrate IRS-1 defines a unique signal transduction protein. *Nature* **352**, 73-77
18. Sun, X. J., Wang, L. M., Zhang, Y., Yenush, L., Myers, M. G., Jr., Glasheen, E., Lane, W. S., Pierce, J. H., and White, M. F. (1995) Role of IRS-2 in insulin and cytokine signalling. *Nature* **377**, 173-177
19. Taniguchi, C. M., Emanuelli, B., and Kahn, C. R. (2006) Critical nodes in signalling pathways: insights into insulin action. *Nat Rev Mol Cell Biol* **7**, 85-96
20. Sun, X. J., Crimmins, D. L., Myers, M. G., Jr., Miralpeix, M., and White, M. F. (1993) Pleiotropic insulin signals are engaged by multisite phosphorylation of IRS-1. *Mol Cell Biol* **13**, 7418-7428
21. Myers, M. G., Jr., Backer, J. M., Sun, X. J., Shoelson, S., Hu, P., Schlessinger, J., Yoakim, M., Schaffhausen, B., and White, M. F. (1992) IRS-1 activates phosphatidylinositol 3'-kinase by associating with src homology 2 domains of p85. *Proc Natl Acad Sci U S A* **89**, 10350-10354
22. Esposito, D. L., Li, Y., Cama, A., and Quon, M. J. (2001) Tyr(612) and Tyr(632) in human insulin receptor substrate-1 are important for full activation of insulin-stimulated phosphatidylinositol 3-kinase activity and translocation of GLUT4 in adipose cells. *Endocrinology* **142**, 2833-2840
23. Chen, H., Nystrom, F. H., Dong, L. Q., Li, Y., Song, S., Liu, F., and Quon, M. J. (2001) Insulin stimulates increased catalytic activity of phosphoinositide-dependent kinase-1 by a phosphorylation-dependent mechanism. *Biochemistry* **40**, 11851-11859
24. Muniyappa, R., Iantorno, M., and Quon, M. J. (2008) An integrated view of insulin resistance and endothelial dysfunction. *Endocrinol Metab Clin North Am* **37**, 685-711, ix-x
25. Dasu, M. R., and Jialal, I. (2011) Free fatty acids in the presence of high glucose amplify monocyte inflammation via Toll-like receptors. *Am J Physiol Endocrinol Metab* **300**, E145-154

26. Akira, S., and Takeda, K. (2004) Toll-like receptor signalling. *Nat Rev Immunol* **4**, 499-511
27. Sun, X. J., and Liu, F. (2009) Phosphorylation of IRS proteins Yin-Yang regulation of insulin signaling. *Vitam Horm* **80**, 351-387
28. Kim, J. A., Yeh, D. C., Ver, M., Li, Y., Carranza, A., Conrads, T. P., Veenstra, T. D., Harrington, M. A., and Quon, M. J. (2005) Phosphorylation of Ser24 in the pleckstrin homology domain of insulin receptor substrate-1 by Mouse Pelle-like kinase/interleukin-1 receptor-associated kinase: cross-talk between inflammatory signaling and insulin signaling that may contribute to insulin resistance. *J Biol Chem* **280**, 23173-23183
29. Hotamisligil, G. S., Peraldi, P., Budavari, A., Ellis, R., White, M. F., and Spiegelman, B. M. (1996) IRS-1-mediated inhibition of insulin receptor tyrosine kinase activity in TNF-alpha- and obesity-induced insulin resistance. *Science* **271**, 665-668
30. Klover, P. J., Zimmers, T. A., Koniaris, L. G., and Mooney, R. A. (2003) Chronic exposure to interleukin-6 causes hepatic insulin resistance in mice. *Diabetes* **52**, 2784-2789
31. Klover, P. J., Clementi, A. H., and Mooney, R. A. (2005) Interleukin-6 depletion selectively improves hepatic insulin action in obesity. *Endocrinology* **146**, 3417-3427
32. Cai, D., Yuan, M., Frantz, D. F., Melendez, P. A., Hansen, L., Lee, J., and Shoelson, S. E. (2005) Local and systemic insulin resistance resulting from hepatic activation of IKK-beta and NF-kappaB. *Nat Med* **11**, 183-190
33. Shoelson, S. E., Herrero, L., and Naaz, A. (2007) Obesity, inflammation, and insulin resistance. *Gastroenterology* **132**, 2169-2180
34. O'Neill, L. A., and Greene, C. (1998) Signal transduction pathways activated by the IL-1 receptor family: ancient signaling machinery in mammals, insects, and plants. *J Leukoc Biol* **63**, 650-657
35. Kenny, E. F., and O'Neill, L. A. (2008) Signalling adaptors used by Toll-like receptors: an update. *Cytokine* **43**, 342-349
36. Jenkins, K. A., and Mansell, A. (2010) TIR-containing adaptors in Toll-like receptor signalling. *Cytokine* **49**, 237-244
37. Li, X., Commane, M., Jiang, Z., and Stark, G. R. (2001) IL-1-induced NFkappa B and c-Jun N-terminal kinase (JNK) activation diverge at IL-1 receptor-associated kinase (IRAK). *Proc Natl Acad Sci U S A* **98**, 4461-4465
38. Wang, Q., Dziarski, R., Kirschning, C. J., Muzio, M., and Gupta, D. (2001) Micrococci and peptidoglycan activate TLR2-->MyD88-->IRAK-->TRAF-->NIK-->IKK-->NF-kappaB signal transduction pathway that induces transcription of interleukin-8. *Infect Immun* **69**, 2270-2276
39. Takaesu, G., Ninomiya-Tsuji, J., Kishida, S., Li, X., Stark, G. R., and Matsumoto, K. (2001) Interleukin-1 (IL-1) receptor-associated kinase leads to activation of TAK1 by inducing TAB2 translocation in the IL-1 signaling pathway. *Mol Cell Biol* **21**, 2475-2484
40. Yuan, M., Konstantopoulos, N., Lee, J., Hansen, L., Li, Z. W., Karin, M., and Shoelson, S. E. (2001) Reversal of obesity- and diet-induced insulin resistance with salicylates or targeted disruption of Ikkbeta. *Science* **293**, 1673-1677
41. Kim, F., Tysseling, K. A., Rice, J., Pham, M., Haji, L., Gallis, B. M., Baas, A. S., Paramsothy, P., Giachelli, C. M., Corson, M. A., and Raines, E. W. (2005) Free fatty acid impairment of nitric oxide production in endothelial cells is mediated by IKKbeta. *Arterioscler Thromb Vasc Biol* **25**, 989-994
42. Kim, F., Pham, M., Maloney, E., Rizzo, N. O., Morton, G. J., Wisse, B. E., Kirk, E. A., Chait, A., and Schwartz, M. W. (2008) Vascular inflammation, insulin resistance, and reduced nitric oxide production precede the onset of peripheral insulin resistance. *Arterioscler Thromb Vasc Biol* **28**, 1982-1988
43. Kim, J. A., Koh, K. K., and Quon, M. J. (2005) The union of vascular and metabolic actions of insulin in sickness and in health. *Arterioscler Thromb Vasc Biol* **25**, 889-891
44. Dewberry, R., Holden, H., Crossman, D., and Francis, S. (2000) Interleukin-1 receptor antagonist expression in human endothelial cells and atherosclerosis. *Arterioscler Thromb Vasc Biol* **20**, 2394-2400
45. McCarty, M. F. (2010) Salsalate may have broad utility in the prevention and treatment of vascular disorders and the metabolic syndrome. *Med Hypotheses* **75**, 276-281
46. Goldfine, A. B., Fonseca, V., Jablonski, K. A., Pyle, L., Staten, M. A., and Shoelson, S. E. (2010) The effects of salsalate on glycemic control in patients with type 2 diabetes: a randomized trial. *Ann Intern Med* **152**, 346-357
47. Fleischman, A., Shoelson, S. E., Bernier, R., and Goldfine, A. B. (2008) Salsalate improves glycemia and inflammatory parameters in obese young adults. *Diabetes Care* **31**, 289-294
48. Lakoski, S. G., Li, L., Langefeld, C. D., Liu, Y., Howard, T. D., Brosnihan, K. B., Xu, J., Bowden, D. W., and Harrington, D. M. (2007) The association between innate immunity gene (IRAK1) and C-reactive protein in the Diabetes Heart Study. *Exp Mol Pathol* **82**, 280-283
49. Bruning, J. C., Michael, M. D., Winnay, J. N., Hayashi, T., Horsch, D., Accili, D., Goodyear, L. J., and Kahn, C. R. (1998) A muscle-specific insulin receptor knockout exhibits features of the metabolic syndrome of NIDDM without altering glucose tolerance. *Mol Cell* **2**, 559-569

50. Michael, M. D., Kulkarni, R. N., Postic, C., Previs, S. F., Shulman, G. I., Magnuson, M. A., and Kahn, C. R. (2000) Loss of insulin signaling in hepatocytes leads to severe insulin resistance and progressive hepatic dysfunction. *Mol Cell* **6**, 87-97
51. Moller, D. E., Chang, P. Y., Yaspekis, B. B., 3rd, Flier, J. S., Wallberg-Henriksson, H., and Ivy, J. L. (1996) Transgenic mice with muscle-specific insulin resistance develop increased adiposity, impaired glucose tolerance, and dyslipidemia. *Endocrinology* **137**, 2397-2405
52. Kido, Y., Burks, D. J., Withers, D., Bruning, J. C., Kahn, C. R., White, M. F., and Accili, D. (2000) Tissue-specific insulin resistance in mice with mutations in the insulin receptor, IRS-1, and IRS-2. *J Clin Invest* **105**, 199-205
53. Ren, J. M., Marshall, B. A., Mueckler, M. M., McCaleb, M., Amatruda, J. M., and Shulman, G. I. (1995) Overexpression of Glut4 protein in muscle increases basal and insulin-stimulated whole body glucose disposal in conscious mice. *J Clin Invest* **95**, 429-432
54. Kim, J. K., Fillmore, J. J., Sunshine, M. J., Albrecht, B., Higashimori, T., Kim, D. W., Liu, Z. X., Soos, T. J., Cline, G. W., O'Brien, W. R., Littman, D. R., and Shulman, G. I. (2004) PKC-theta knockout mice are protected from fat-induced insulin resistance. *J Clin Invest* **114**, 823-827
55. Chen, H., Sullivan, G., and Quon, M. J. (2005) Assessing the predictive accuracy of QUICKI as a surrogate index for insulin sensitivity using a calibration model. *Diabetes* **54**, 1914-1925
56. Quon, M. J. (2002) QUICKI is a useful and accurate index of insulin sensitivity. *J Clin Endocrinol Metab* **87**, 949-951
57. Hotamisligil, G. S. (2000) Molecular mechanisms of insulin resistance and the role of the adipocyte. *Int J Obes Relat Metab Disord* **24 Suppl 4**, S23-27
58. Uysal, K. T., Wiesbrock, S. M., Marino, M. W., and Hotamisligil, G. S. (1997) Protection from obesity-induced insulin resistance in mice lacking TNF-alpha function. *Nature* **389**, 610-614
59. Jiang, Z. Y., Zhou, Q. L., Chatterjee, A., Feener, E. P., Myers, M. G., Jr., White, M. F., and King, G. L. (1999) Endothelin-1 modulates insulin signaling through phosphatidylinositol 3-kinase pathway in vascular smooth muscle cells. *Diabetes* **48**, 1120-1130
60. Potenza, M. A., Marasciulo, F. L., Chieppa, D. M., Brigiani, G. S., Formoso, G., Quon, M. J., and Montagnani, M. (2005) Insulin resistance in spontaneously hypertensive rats is associated with endothelial dysfunction characterized by imbalance between NO and ET-1 production. *Am J Physiol Heart Circ Physiol* **289**, H813-822
61. Kim, J. A., Montagnani, M., Koh, K. K., and Quon, M. J. (2006) Reciprocal relationships between insulin resistance and endothelial dysfunction: molecular and pathophysiological mechanisms. *Circulation* **113**, 1888-1904
62. Ayala, J. E., Samuel, V. T., Morton, G. J., Obici, S., Croniger, C. M., Shulman, G. I., Wasserman, D. H., McGuinness, O. P., and Consortium, N. I. H. M. M. P. C. (2010) Standard operating procedures for describing and performing metabolic tests of glucose homeostasis in mice. *Dis Model Mech* **3**, 525-534
63. Franceschi, C., Bonafe, M., Valensin, S., Olivieri, F., De Luca, M., Ottaviani, E., and De Benedictis, G. (2000) Inflamm-aging. An evolutionary perspective on immunosenescence. *Ann NY Acad Sci* **908**, 244-254
64. Minciullo, P. L., Catalano, A., Mandraffino, G., Casciaro, M., Crucitti, A., Maltese, G., Morabito, N., Lasco, A., Gangemi, S., and Basile, G. (2016) Inflammaging and Anti-Inflammaging: The Role of Cytokines in Extreme Longevity. *Arch Immunol Ther Exp (Warsz)* **64**, 111-126
65. Jang, H. J., Kim, H. S., Hwang, D. H., Quon, M. J., and Kim, J. A. (2013) Toll-like receptor 2 mediates high-fat diet-induced impairment of vasodilator actions of insulin. *Am J Physiol Endocrinol Metab* **304**, E1077-1088
66. Berglund, M., Thomas, J. A., Hornquist, E. H., and Hultgren, O. H. (2008) Toll-like receptor cross-hyporesponsiveness is functional in interleukin-1-receptor-associated kinase-1 (IRAK-1)-deficient macrophages: differential role played by IRAK-1 in regulation of tumour necrosis factor and interleukin-10 production. *Scand J Immunol* **67**, 473-479
67. Swantek, J. L., Tsen, M. F., Cobb, M. H., and Thomas, J. A. (2000) IL-1 receptor-associated kinase modulates host responsiveness to endotoxin. *J Immunol* **164**, 4301-4306
68. Uematsu, S., Sato, S., Yamamoto, M., Hirotani, T., Kato, H., Takeshita, F., Matsuda, M., Coban, C., Ishii, K. J., Kawai, T., Takeuchi, O., and Akira, S. (2005) Interleukin-1 receptor-associated kinase-1 plays an essential role for Toll-like receptor (TLR)7- and TLR9-mediated interferon- induction. *J Exp Med* **201**, 915-923
69. Maitra, U., Chang, S., Singh, N., and Li, L. (2009) Molecular mechanism underlying the suppression of lipid oxidation during endotoxemia. *Mol Immunol* **47**, 420-425
70. Van Heek, M., Compton, D. S., France, C. F., Tedesco, R. P., Fawzi, A. B., Graziano, M. P., Sybertz, E. J., Strader, C. D., and Davis, H. R., Jr. (1997) Diet-induced obese mice develop peripheral, but not central, resistance to leptin. *J Clin Invest* **99**, 385-390
71. Sell, H., Habich, C., and Eckel, J. (2012) Adaptive immunity in obesity and insulin resistance. *Nat Rev Endocrinol* **8**, 709-716

72. Kim, Y. B., Shulman, G. I., and Kahn, B. B. (2002) Fatty acid infusion selectively impairs insulin action on Akt1 and protein kinase C lambda /zeta but not on glycogen synthase kinase-3. *J Biol Chem* **277**, 32915-32922
73. Yu, C., Chen, Y., Cline, G. W., Zhang, D., Zong, H., Wang, Y., Bergeron, R., Kim, J. K., Cushman, S. W., Cooney, G. J., Atcheson, B., White, M. F., Kraegen, E. W., and Shulman, G. I. (2002) Mechanism by which fatty acids inhibit insulin activation of insulin receptor substrate-1 (IRS-1)-associated phosphatidylinositol 3-kinase activity in muscle. *J Biol Chem* **277**, 50230-50236
74. Eckardt, K., Taube, A., and Eckel, J. (2011) Obesity-associated insulin resistance in skeletal muscle: role of lipid accumulation and physical inactivity. *Rev Endocr Metab Disord* **12**, 163-172
75. Mishra, A. K., Dubey, V., and Ghosh, A. R. (2016) Obesity: An overview of possible role(s) of gut hormones, lipid sensing and gut microbiota. *Metabolism* **65**, 48-65
76. Talior, I., Yarkoni, M., Bashan, N., and Eldar-Finkelman, H. (2003) Increased glucose uptake promotes oxidative stress and PKC-delta activation in adipocytes of obese, insulin-resistant mice. *Am J Physiol Endocrinol Metab* **285**, E295-302
77. Kim, O. K., Jun, W., and Lee, J. (2015) Mechanism of ER Stress and Inflammation for Hepatic Insulin Resistance in Obesity. *Ann Nutr Metab* **67**, 218-227
78. Lambert, G. W., Straznicky, N. E., Lambert, E. A., Dixon, J. B., and Schlaich, M. P. (2010) Sympathetic nervous activation in obesity and the metabolic syndrome--causes, consequences and therapeutic implications. *Pharmacol Ther* **126**, 159-172
79. Esteve, E., Ricart, W., and Fernandez-Real, J. M. (2011) Gut microbiota interactions with obesity, insulin resistance and type 2 diabetes: did gut microbiota co-evolve with insulin resistance? *Curr Opin Clin Nutr Metab Care* **14**, 483-490
80. Nawaratne, R., Gray, A., Jorgensen, C. H., Downes, C. P., Siddle, K., and Sethi, J. K. (2006) Regulation of insulin receptor substrate 1 pleckstrin homology domain by protein kinase C: role of serine 24 phosphorylation. *Mol Endocrinol* **20**, 1838-1852
81. Greene, M. W., Ruhoff, M. S., Roth, R. A., Kim, J. A., Quon, M. J., and Krause, J. A. (2006) PKCdelta-mediated IRS-1 Ser24 phosphorylation negatively regulates IRS-1 function. *Biochem Biophys Res Commun* **349**, 976-986
82. Han, S. H., Quon, M. J., and Koh, K. K. (2007) Reciprocal relationships between abnormal metabolic parameters and endothelial dysfunction. *Curr Opin Lipidol* **18**, 58-65
83. Chen, H., Lin, A. S., Li, Y., Reiter, C. E., Ver, M. R., and Quon, M. J. (2008) Dehydroepiandrosterone stimulates phosphorylation of FoxO1 in vascular endothelial cells via phosphatidylinositol 3-kinase- and protein kinase A-dependent signaling pathways to regulate ET-1 synthesis and secretion. *J Biol Chem* **283**, 29228-29238
84. Formoso, G., Chen, H., Kim, J. A., Montagnani, M., Consoli, A., and Quon, M. J. (2006) Dehydroepiandrosterone mimics acute actions of insulin to stimulate production of both nitric oxide and endothelin 1 via distinct phosphatidylinositol 3-kinase- and mitogen-activated protein kinase-dependent pathways in vascular endothelium. *Mol Endocrinol* **20**, 1153-1163
85. Muniyappa, R., Montagnani, M., Koh, K. K., and Quon, M. J. (2007) Cardiovascular actions of insulin. *Endocr Rev* **28**, 463-491
86. Rizza, R. A., Haymond, M. W., Verdonk, C. A., Mandarino, L. J., Miles, J. M., Service, F. J., and Gerich, J. E. (1981) Pathogenesis of hypoglycemia in insulinoma patients: suppression of hepatic glucose production by insulin. *Diabetes* **30**, 377-381
87. Nishimura, H., Pallardo, F. V., Seidner, G. A., Vannucci, S., Simpson, I. A., and Birnbaum, M. J. (1993) Kinetics of GLUT1 and GLUT4 glucose transporters expressed in *Xenopus* oocytes. *J Biol Chem* **268**, 8514-8520
88. Lee, S., Lynn, E. G., Kim, J. A., and Quon, M. J. (2008) Protein kinase C-zeta phosphorylates insulin receptor substrate-1, -3, and -4 but not -2: isoform specific determinants of specificity in insulin signaling. *Endocrinology* **149**, 2451-2458
89. Cong, L. N., Chen, H., Li, Y., Zhou, L., McGibbon, M. A., Taylor, S. I., and Quon, M. J. (1997) Physiological role of Akt in insulin-stimulated translocation of GLUT4 in transfected rat adipose cells. *Mol Endocrinol* **11**, 1881-1890
90. Fasshauer, M., Klein, J., Ueki, K., Kriauciunas, K. M., Benito, M., White, M. F., and Kahn, C. R. (2000) Essential role of insulin receptor substrate-2 in insulin stimulation of Glut4 translocation and glucose uptake in brown adipocytes. *J Biol Chem* **275**, 25494-25501
91. Toye, A. A., Lippiat, J. D., Proks, P., Shimomura, K., Bentley, L., Hugill, A., Mijat, V., Goldsworthy, M., Moir, L., Haynes, A., Quarterman, J., Freeman, H. C., Ashcroft, F. M., and Cox, R. D. (2005) A genetic and physiological study of impaired glucose homeostasis control in C57BL/6J mice. *Diabetologia* **48**, 675-686
92. Freeman, H., Shimomura, K., Horner, E., Cox, R. D., and Ashcroft, F. M. (2006) Nicotinamide nucleotide transhydrogenase: a key role in insulin secretion. *Cell Metab* **3**, 35-45

93. Sperry, J. L., Zolin, S., Zuckerbraun, B. S., Vodovotz, Y., Namas, R., Neal, M. D., Ferrell, R. E., Rosengart, M. R., Peitzman, A. B., and Billiar, T. R. (2014) X chromosome-linked IRAK-1 polymorphism is a strong predictor of multiple organ failure and mortality postinjury. *Ann Surg* **260**, 698-703; discussion 703-695
94. He, L., Sabet, A., Djedjos, S., Miller, R., Sun, X., Hussain, M. A., Radovick, S., and Wondisford, F. E. (2009) Metformin and insulin suppress hepatic gluconeogenesis through phosphorylation of CREB binding protein. *Cell* **137**, 635-646
95. Folli, F., Saad, M. J., Backer, J. M., and Kahn, C. R. (1992) Insulin stimulation of phosphatidylinositol 3-kinase activity and association with insulin receptor substrate 1 in liver and muscle of the intact rat. *J Biol Chem* **267**, 22171-22177
96. Saad, M. J., Araki, E., Miralpeix, M., Rothenberg, P. L., White, M. F., and Kahn, C. R. (1992) Regulation of insulin receptor substrate-1 in liver and muscle of animal models of insulin resistance. *J Clin Invest* **90**, 1839-1849
97. Saad, M. J., Folli, F., Kahn, J. A., and Kahn, C. R. (1993) Modulation of insulin receptor, insulin receptor substrate-1, and phosphatidylinositol 3-kinase in liver and muscle of dexamethasone-treated rats. *J Clin Invest* **92**, 2065-2072
98. Mirbolooki, M. R., Upadhyay, S. K., Constantinescu, C. C., Pan, M. L., and Mukherjee, J. (2014) Adrenergic pathway activation enhances brown adipose tissue metabolism: a [(1)(8)F]FDG PET/CT study in mice. *Nucl Med Biol* **41**, 10-16
99. Thorn, S. L., deKemp, R. A., Dumouchel, T., Klein, R., Renaud, J. M., Wells, R. G., Gollob, M. H., Beanlands, R. S., and DaSilva, J. N. (2013) Repeatable noninvasive measurement of mouse myocardial glucose uptake with 18F-FDG: evaluation of tracer kinetics in a type 1 diabetes model. *J Nucl Med* **54**, 1637-1644
100. Chen, H., Sullivan, G., Yue, L. Q., Katz, A., and Quon, M. J. (2003) QUICKI is a useful index of insulin sensitivity in subjects with hypertension. *Am J Physiol Endocrinol Metab* **284**, E804-812
101. Burant, C. F., and Bell, G. I. (1992) Mammalian facilitative glucose transporters: evidence for similar substrate recognition sites in functionally monomeric proteins. *Biochemistry* **31**, 10414-10420

FIGURE LEGNDS

Figure 1. Intraperitoneal glucose tolerance tests (IPGTT): 6-week old wild type and *Irak1* k/o mice were fed with either low fat (panels A and B, n = 8) or high fat diets (panel C and D, n = 8). After 12 weeks on the designated diets, mice were fasted for 5 h. Glucose (1 g/kg) was injected intraperitoneally and blood glucose was measured at 0, 10, 20, 30, 60 and 120 min after injection. Areas under the curve (AUC) were calculated (panels B and D) based on the trapezoidal rule. **Panel A:** ANOVA for repeated measures, $p = 0.004$. **Panel B:** $AUC_{\text{glucose}} = 10890 \pm 840$ vs. 6850 ± 1100 (mg/dl)•min, $p = 0.035$. **Panel C:** ANOVA for repeated measures, $p = 0.56$; **Panel D:** $AUC_{\text{glucose}} = 14370 \pm 1800$ vs. 15180 ± 1040 (mg/dl)•min, $p = 0.59$. Data are expressed as mean \pm SEM.

Figure 2. Hyperinsulinemic euglycemic glucose clamp studies in mice fed normal chow diet. At 28 weeks of age, glucose clamp studies were performed as described in Experimental Procedures, and previously (94). Mean glucose infusion rates over the 30 min steady-state period at the end of the study were used to estimate the steady-state glucose infusion rate (GIR). Data shown are mean \pm SEM. **Panel A:** For the WT mice (n = 8), blood glucose is indicated with solid circles (●) and GIR with closed triangles (▲). **Panel B:** For *Irak1* k/o mice (n = 10), blood glucose is indicated with open circles (○) and GIR with open triangles (Δ). **Panel C:** GIR normalized for body weight. **Panel D:** Insulin Sensitivity Index (ISI) was defined as the GIR (glucose mg/kg/min) divided by the mean glucose level (mg/dl) calculated from the last 30 min of the clamp study when steady-state had been achieved.

Figure 3. Cardiometabolic biomarkers in mice fed with low fat diet (A – F) and high fat diet (G – L). Fasting plasma was obtained after 14 weeks on low fat diet or high fat diet. Glucose (Panel B and H) was measured by YSI 2700 Select Biochemistry Analyzer; insulin (Panel A and G), adiponectin (Panel D and J), ET-1 (Panel E and K) and CRP (Panel F and L) were measured by ELISA. Quantitative Insulin Sensitivity Check Index (QUICKI) (Panel C and I) was calculated as: $QUICKI = 1 / [\log(\text{insulin}) + \log(\text{glucose})]$ (55,56,100). Data are expressed as mean \pm SEM.

Figure 4. *In vivo* insulin-stimulated [^{18}F]fluorodeoxyglucose (^{18}F -FDG) distribution in mice fed with normal chow diet. After mice were fasted overnight, ^{18}F -FDG 11.1 MBq (300 Ci) and insulin (1.125U/kg) in 200 μL of saline were injected via tail vein under anesthesia. Dynamic imaging was acquired for 30 min by Siemens Inveon Small Animal PET-CT Imaging System. The data were obtained by drawing Regions of Interest (ROI) over target organs of muscle, abdominal subcutaneous white adipose tissue, liver, and brain (98,99). The ^{18}F -FDG bio-distribution in the target organs/tissues were analyzed at three time points: 10, 20 and 30 minutes. **Panel A:** Time-dependent distribution [% of injected dose (ID) per gram] of ^{18}F -FDG in the hind limb, liver, and abdominal white fat in *Irak1* k/o mice and in wild-type mice. **Panel B:** Relative time-dependent distribution of ^{18}F -FDG in the hind limb, liver, and abdominal white fat (ratio of ^{18}F -FDG: specific tissue/brain) in *Irak1* k/o mice and in wild-type mice. WT n = 3; k/o n = 3, * $p < 0.04$, ** $p < 0.03$. Data shown are mean \pm SEM.

Figure 5. Hyperinsulinemic euglycemic glucose clamp studies with [$3\text{-}^3\text{H}$]glucose tracer in mice fed with a normal chow diet. 6 WT and 6 k/o mice underwent glucose clamps with ^3H -glucose tracer as described in Experimental Procedures, and previously (94). At the beginning and the end of each clamp study, blood was collected for calculating insulin-stimulated glucose disposal rates and hepatic glucose production rate. **Panel A:** GIR normalized for body weight. **Panel B:** ISI was defined as GIR (mg/kg/min) divided by steady-state glucose level (mg/dl) during the last 30 min of the clamp. **Panel C:** Glucose disposal rate. **Panel D:** Hepatic glucose production rate. Data are expressed as mean \pm SEM.

Figure 6. Immunoblotting analysis of insulin signaling pathways in muscle (panels A-B), liver (panels C-D), and abdominal white adipose tissue (panels E-F). 20 weeks old mice fed with chow diet were fasted for 5 h and then given saline (None) or insulin (10 U/kg body weight) via portal vein injection under anesthesia. 5 min after injection, skeletal muscle from hind limb, abdominal fat, and liver were immediately flash frozen and homogenized. These samples were centrifuged and supernatants of tissue lysates were

analyzed by immunoblotting with indicated antibodies. Immunoblots with samples from 6 WT and 7 *Irak1* k/o mice are shown in panels A, C and E. Immunoblots for phospho-Akt (panels B, D, and F) were quantified by scanning densitometry and normalized for Akt total protein. Data are expressed as mean \pm SEM.

FIGURES

Figure 1




Figure 2




Figure 3




Figure 4




Figure 5





Figure 6



Deletion of interleukin 1 receptor associated kinase 1 (*IRAK1*) improves glucose tolerance primarily by increasing insulin sensitivity in skeletal muscle

Xiao Jian Sun, Soohyun Park Kim, Dongming Zhang, Helen Sun, Qi Cao, Xin Lu, Zhekang Ying, Liwu Li, Robert R. Henry, Theodore P. Ciaraldi, Simeon I. Taylor and Michael J. Quon

J. Biol. Chem. published online June 1, 2017

Access the most updated version of this article at doi: [10.1074/jbc.M117.779108](https://doi.org/10.1074/jbc.M117.779108)

Alerts:

- [When this article is cited](#)
- [When a correction for this article is posted](#)

[Click here](#) to choose from all of JBC's e-mail alerts

Supplemental material:

<http://www.jbc.org/content/suppl/2017/06/01/M117.779108.DC1>

This article cites 0 references, 0 of which can be accessed free at

<http://www.jbc.org/content/early/2017/06/01/jbc.M117.779108.full.html#ref-list-1>