

**THE ACTIVITIES OF THE C-TERMINAL REGIONS OF THE FORMIN PROTEIN
DISHEVELED-ASSOCIATED ACTIVATOR OF MORPHOGENESIS (DAAM) IN ACTIN
DYNAMICS**

Andrea Teréz Vig¹, István Földi², Szilárd Szikora², Ede Migh², Rita Gombos², Mónika Ágnes Tóth¹,
Tamás Huber¹, Réka Pintér¹, Gábor Csaba Talián¹, József Mihály², Beáta Bugyi^{1,3,*}

* to whom the correspondence should be addressed: Beáta Bugyi, University of Pécs, Medical School,
Department of Biophysics, Pécs, Szigeti str. 12, H-7624, Hungary, Phone: +36-72-536-265, E-mail:
beata.bugyi@aok.pte.hu

Institutions:

¹ University of Pécs, Medical School, Department of Biophysics, Szigeti str. 12, Pécs, H-7624, Hungary

² Biological Research Centre, Hungarian Academy of Sciences, Institute of Genetics, MTA-SZBK NAP
B Axon Growth and Regeneration Group, Temesvári krt. 62, Szeged, H-6726, Hungary

³ Szentágotthai Research Center, Ifjúság str. 34, Pécs, H-7624, Hungary

Running title:

The C-terminus of formins in actin dynamics

Keywords:

actin, formin, *Drosophila*, neuron, fluorescence

ABSTRACT

Disheveled-associated activator of morphogenesis (DAAM) is a diaphanous-related formin protein essential for the regulation of actin cytoskeleton dynamics in diverse biological processes. The conserved formin homology 1 and 2 (FH1–FH2) domains of DAAM catalyze actin nucleation and processively mediate filament elongation. These activities are indirectly regulated by the N-, and C-terminal regions flanking the FH1–FH2 domains. Recently, the C-terminal diaphanous-autoregulatory domain (DAD) and the C-terminus (CT) of formins have also been shown to regulate actin assembly by directly interacting with actin. Here, to better understand the biological activities of DAAM, we studied the role of DAD–CT regions of *Drosophila* DAAM in its interaction with actin with *in vitro* biochemical and *in vivo* genetic approaches. We found that the DAD–CT region binds actin *in vitro* and that its main actin-binding element is the CT region, which does not influence actin dynamics on its own. However, we also found that it can tune the nucleating activity and the filament end–interaction properties of DAAM in an FH2 domain–dependent manner. We also demonstrate that DAD-CT makes the FH2 domain more efficient in antagonizing with capping protein. Consistently, *in vivo* data suggested that the CT region contributes to DAAM-mediated filopodia formation and dynamics in primary neurons. In conclusion, our results demonstrate that the CT region of DAAM plays an important role in actin assembly regulation in a biological context.

INTRODUCTION

Formins are actin assembly machineries playing essential roles in diverse biological processes. The core actin-interacting region is the conserved Formin homology 2 (FH2)¹ domain that catalyzes actin nucleation and mediates processive elongation of filament ends. The activities of the FH2 domain are assisted by the upstream Formin homology 1 (FH1) domain that interacts with profilin-actin (reviewed in (1-5)). The N-, and C-terminal regions flanking the FH1-FH2 domains have diverse composition and functions amongst different formin families. Formins belonging to the Diaphanous-related formin (DRF) family including Diaphanous (Dia), Disheveled-associated activator of morphogenesis (DAAM), Formin-like protein (FMNL), FH1/FH2 domain-containing protein (FHOD) and Inverted formin2 (INF2) are characterized by the presence of an N-terminal Diaphanous inhibitory domain (DID) and a C-terminal Diaphanous autoregulatory domain (DAD). It is well established that the intramolecular contacts formed between these regions keep the FH2 in an inactive state by preventing its interaction with actin. The functions of the FH2 domain in actin assembly are activated upon Rho GTPase-dependent relief of the DID:DAD autoinhibitory interaction (5-11).

Recent studies showed that, besides autoregulation, the C-terminal regions of formins from yeast to mammals (such as mouse Dia1, FMNL3, INF2, *Drosophila* Capuccino, yeast Bni1 and Bnr1 and human Daam1) can also influence the active FH2 domain mediated actin assembly (12-15). A common feature of the C-terminal regions studied so far is that they increase the efficiency of the FH2-catalyzed nucleation. Additionally, isolated C-terminal regions can directly interact with actin, independently of the FH2 domain. The C-terminus of INF2 contains a Wiskott-Aldrich syndrome homology (WH2)-like DAD motif, which in its isolated form sequesters monomeric actin and severs actin filaments (12), while the WH2-DAD-CT region of

FMNL3 nucleates actin and slows elongation in a dimeric form (14). Similarly, the isolated dimeric DAD from mDia1 seems to be sufficient to promote actin nucleation (13). In contrast, the tail region of Capuccino does not influence nucleation or elongation in the absence of FH2 (15). The biochemical evidences suggest that the activities of the C-terminal regions vary amongst formins, and raise the question how and which of the activities of the isolated C-terminal regions are transmitted to the functionality of each formin in the context of the FH1-FH2 domains. In addition, the *in vivo* significance of the direct interaction of the C-terminal regions of formins with actin and its role in actin cytoskeleton dynamics regulation is not well understood.

In this work, we investigated the *Drosophila* DAAM formin belonging to the DRF family. DAAM is involved in diverse morphogenetic processes mediated by the actin cytoskeleton. For example, DAAM plays a role in organizing apical actin cables that define the tracheal cuticle pattern (16). It is also required for axonal growth and guidance by promoting filopodia formation in the growth cone (17), and DAAM is essential for sarcomerogenesis (18-20). In our previous work we described the physico-chemical properties of the interaction of *Drosophila* DAAM FH1-FH2 and actin, and showed that DAAM FH1-FH2 is a profilin-gated actin assembly factor (18,21). To better understand the biological functioning of DAAM, we aimed to analyze the biochemical activities of recombinantly produced DAD-CT regions of DAAM in its isolated form, as well as in the context of the FH1-FH2 domains in the regulation of actin assembly. By dissecting the activities of the DAD and CT regions of DAAM we aimed to reveal which of the reported activities of the C-terminal regions of formins are shared by the DAD-CT regions of *Drosophila* DAAM to influence FH1-FH2-assisted actin assembly. We also investigated how the DAD-CT region influences filopodia formation and FH2-mediated actin dynamics in primary neurons using *in vivo* genetic approaches.

¹ Abbreviations:

Formin homology (FH), Diaphanous-related formin family (DRF), Diaphanous-related formin (Dia), Disheveled-associated activator of morphogenesis (DAAM), Formin-like protein (FMNL), FH1/FH2

domain-containing protein (FHOD), Inverted formin (INF2), Diaphanous inhibitory domain (DID), Diaphanous autoregulatory domain (DAD), Wiskott-Aldrich syndrome homology (WH2), Total internal reflection fluorescence microscopy (TIRFM), Capping Protein (CP)

RESULTS

cDAAM is more efficient in catalyzing actin nucleation than the FH1-FH2 domains

To investigate the effects of the C-terminal regions of *Drosophila* DAAM on actin assembly, first we produced two recombinant proteins that either lack or possess the C-terminal DAD-CT regions, the FH1-FH2 and cDAAM proteins, respectively (Figure 1, A). To analyze the actin assembly efficiencies of these constructs, the polymerization kinetics of G-actin and profilin:G-actin were measured in pyrenyl polymerization experiments (Figure 2, A and B). We found that both constructs can accelerate actin polymerization (Figure 2, A). However, cDAAM accelerates the overall polymerization rate of both free G-actin and profilin:G-actin ~ 36fold more efficiently than the FH1-FH2 (which lacks the C-terminal DAD-CT regions) at all concentrations that were tested (Figure 2, B). This observation is consistent with the findings on the human Daam1 protein, suggesting that this feature is conserved between species (13).

Since the FH2 domain of some formins, such as Bni1 and mDia1, has low affinity to monomeric actin in the absence of the C-terminal regions (13,22,23), we wanted to examine whether the differences in the actin assembly promoting activities of FH1-FH2 and cDAAM can arise from different actin affinities. Therefore, the interactions of FH1-FH2 and cDAAM with G-actin and profilin:G-actin were investigated in steady-state fluorescence anisotropy experiments. To avoid artificial increase in anisotropy due to the presence of polymerizing agents, LatrunculinA was used to keep actin in monomeric form (24,25). We observed a similar concentration dependent increase in the values of the steady-state anisotropy in the presence of both DAAM constructs (Figure 2, C). This reflects that FH1-FH2, as well as cDAAM binds monomeric actin and the complexes are characterized by similar affinities in the few tens of nM range (Table 1). This result suggests that the lower polymerization activity of FH1-FH2 compared to cDAAM does not originate from their different affinities to monomeric actin.

The different polymerization efficiencies of FH1-FH2 and cDAAM can also arise from differences in their nucleation and/or elongation activities. Since bulk pyrenyl

polymerization assays cannot distinguish these two activities, the assembly of individual actin filaments was investigated by total internal reflection fluorescence microscopy (TIRFM) (Figure 2, D - F). In control experiments we found that the number of actin filaments significantly decreased in the presence of profilin, compared to that of measured for free G-actin ($p < 0.0001$, Figure 2, D and E), consistent with the nucleation suppressing activity of profilin (26,27). In the presence of FH1-FH2 and cDAAM a dramatic increase in the number of filaments formed was observed as compared to the number of spontaneously formed filaments, which reflects the nucleation promoting activity of these constructs (Figure 2, D and E). Importantly, quantitative analysis of these data revealed that cDAAM nucleated significantly more filaments than FH1-FH2, suggesting that cDAAM is more efficient in promoting the initial nucleation phase of actin filament formation than FH1-FH2 (Figure 2, D and E). This result demonstrates that the DAD-CT region makes the FH2 domain more efficient in catalyzing actin nucleation. Spontaneously growing actin filaments elongated at a rate of 4.30 ± 0.6 subunit \times s $^{-1}$ and 3.48 ± 0.4 subunit \times s $^{-1}$ in the absence and presence of profilin, respectively (Figure 2, F). These values are in agreement with the well-established barbed end association rate constants of ATP-Mg $^{2+}$ -G-actin (free G-actin: $k_+ = 10.23 \pm 1.5$ μ M $^{-1}\times$ s $^{-1}$, profilin:G-actin: $k_+ = 8.29 \pm 0.9$ μ M $^{-1}\times$ s $^{-1}$ Eq. 4. (21,28,29)). The analysis showed that both FH1-FH2 and cDAAM almost completely inhibit the elongation of free G-actin ($v_{\text{FH1-FH2}} = 0.36 \pm 0.2$ subunit \times s $^{-1}$ ($p < 0.0001$), $v_{\text{cDAAM}} = 0.37 \pm 0.1$ subunit \times s $^{-1}$ ($p < 0.0001$), Figure 2, F). In contrast, in the presence of profilin the elongation rate is similar to that of the spontaneously growing actin filaments ($v_{\text{FH1-FH2}} = 3.44 \pm 0.3$ subunit \times s $^{-1}$ ($p = 0.45$), $v_{\text{cDAAM}} = 3.49 \pm 0.4$ subunit \times s $^{-1}$ ($p = 0.44$), Figure 2, F). Thus, profilin by relieving the elongation inhibitory activity of the constructs contributes to effective filament growth. These observations, regarding the profilin-gated effect of FH1-FH2 on filament elongation are entirely consistent with our previously published data (21). The profilin-dependent regulation of filament elongation was also reported for the human Daam1 protein (30), however the net effect was different. Daam1 did not affect filament growth significantly in the absence of

profilin, while it accelerated elongation from profilin-actin ~ 4 fold as compared to the spontaneous rate (30). These differences may arise from the different conditions including the species specific differences in activities, the expression system, the tag, the profilin isoform used and/or the labeling of hDaam1.

In conclusion, the effects of the FH1-FH2 and cDAAM constructs on filament elongation from either free, or profilin:G-actin are quantitatively the same, however their nucleation efficiencies differ markedly. These observations clearly show that the more potent actin assembly activity of cDAAM as compared to FH1-FH2 arises from a difference in their nucleating efficiency, suggesting that the C-terminal regions contribute to the nucleating activity of the FH2 domain.

The main G-actin binding site in DAD-CT is the CT region

To analyze the activities of the C-terminal regions of DAAM in more details, recombinant fragments were produced containing both the DAD domain and the C-terminal end of the protein (DAD-CT) or only the isolated DAD domain (DAD) (*Figure 1, A*). First, the interaction of DAD-CT and DAD with monomeric actin was investigated by measuring the steady-state anisotropy of fluorescently labelled actin. We found that DAD-CT can bind to G-actin, independently of the FH2 domain (*Figure 3, A*). However, the increase in salt concentration (i.e. increase in the amount of NaCl) resulted in a gradual decrease in actin affinity, suggesting an ionic strength-dependent interaction (*Figure 3, A, Table 1*), which is similar to the salt-dependent interaction reported for the tail domain of Capuccino (15). DAD-CT interacts with profilin:G-actin equally well as with free G-actin indicating that profilin does not affect the binding (*Figure 3, A, Table 1*). In contrast to DAD-CT, the interaction of the DAD domain lacking the CT region with monomeric actin is extremely weak even at low ionic strength conditions ($K_D > 100 \mu\text{M}$ at 5 mM NaCl) (*Figure 3, A, Table 1*). To further analyze which regions are important in the interaction between DAD-CT and monomeric actin, we investigated a construct in which the basic arginine residues found in the CT region were replaced by alanines (DAD-CT^{R-A}) (*Figure 1, A and C*). Our data did not show a detectable interaction between DAD-CT^{R-A} and

monomeric actin ($K_D > 500 \mu\text{M}$ at 5 mM NaCl) (*Figure 3, A, Table 1*).

To further elaborate on the binding mode of DAD-CT, the influence of different actin-binding proteins was tested on its interaction with G-actin. Profilin is known to interact with the hydrophobic cleft of monomeric actin (31). The steady-state anisotropy of fluorescently labelled profilin in complex with G-actin was measured in the presence of DAD-CT. We found that DAD-CT causes a concentration dependent increase in the steady-state anisotropy of profilin bound to G-actin (*Figure 3, B, Table 1*). In case of competitive binding, one would expect the dissociation of fluorescent profilin from G-actin, which would result in a decrease in anisotropy. The opposite tendency that we observed suggests that a ternary complex is formed between monomeric actin, profilin and DAD-CT, indicating that the core binding site of DAD-CT is different from that of profilin. Next, we tested how the interaction of DAD-CT is influenced by WH2 domain proteins, which N-terminally interact with the barbed face, while their C-terminal binds along the ridge between the inner and outer domains of actin up to the pointed end (32-34). We used the WH2 domains of Sarcomere length short protein (SALS-WH2), which have relatively long C-terminal extensions linked to the N-terminal α -helix (28). The steady-state anisotropy of the DAD-CT:G-actin complex was decreased upon addition of SALS-WH2 in a concentration dependent manner (*Figure 3, C*). At high SALS-WH2 concentration the value of the anisotropy decreased to the value characteristic to the SALS-WH2:G-actin complex, suggesting that SALS-WH2 interferes with the G-actin binding of DAD-CT. This indicates that the main binding site of DAD-CT overlaps with that of WH2.

Altogether, these observations support that the isolated C-terminus of DAAM can interact with monomeric actin, and that the core C-terminal actin-binding elements are located mainly in the CT region that is able to establish electrostatic interactions with actin.

DAD-CT does not influence actin dynamics in the absence of the FH2 domain

To test the functional consequences of monomer binding, the effects of isolated DAD-CT on actin dynamics were investigated. In

fluorescence spectroscopy experiments we found that DAD-CT does not affect significantly the assembly of free G-actin and profilin:G-actin at lower concentrations (Figure 3, D). However, higher amounts of DAD-CT ($> \sim 40 - 50 \mu\text{M}$) resulted in a concentration dependent decrease in the pyrenyl fluorescence, similarly to what was observed for Capuccino tail (15). In agreement with the spectroscopic data, TIRFM experiments showed that DAD-CT cannot affect significantly actin nucleation (Figure 3, E and F) and it does not affect the rate of elongation at lower concentrations either (*data not shown*). DAD-CT moderately slows elongation of free G-actin when it is present at higher amounts ($> \sim 40 \mu\text{M}$) ($v_{\text{DAD-CT}} = 3.79 \pm 0.4 \text{ subunit} \times \text{s}^{-1}$ ($p = 0.009$), Figure 3, G), but it does not influence elongation from profilin:G-actin ($v_{\text{DAD-CT}} = 3.64 \pm 0.6 \text{ subunit} \times \text{s}^{-1}$ ($p = 0.47$), Figure 3, G). Also, DAD and DAD-CT^{R-A} do not show any effect in the above experiments, consistently with their negligible interaction with monomeric actin (Figure 3, G).

In conclusion, in contrast to the DAD region of mDial (13), the isolated DAAM DAD-CT cannot influence significantly actin assembly in the absence of the FH2 domain, which suggests that the C-terminal region of DAAM possesses an FH2-dependent function in the regulation of actin dynamics.

Functional CT is required for the full nucleation promoting activity of DAAM

To further elaborate on the contribution of the DAD-CT region to the nucleation activity of the FH2 domain, we tested its effect in the presence of the FH2 domain. For this purpose, two C-terminally modified constructs were generated possessing the native FH2 but lacking a functional CT region; either by deleting the CT region (cDAAM Δ CT) or by introducing the R-A mutation into the CT region (cDAAM^{R-A}) (Figure 1, A). The effects of the two constructs are indistinguishable from each other both in pyrenyl and TIRFM actin assembly assays, which supports that the two modifications are equivalent (Figure 4). However, neither cDAAM Δ CT nor cDAAM^{R-A} can retrieve the maximum nucleation promoting activity of cDAAM. Both C-terminally modified constructs are ~ 6 fold more efficient in accelerating the overall rate of actin polymerization than FH1-FH2, but ~ 6 fold less effective than cDAAM possessing the native

CT region (Figure 4, A and B). Since they influence elongation in a similar manner as the wild type constructs, the polymerization promoting effect of cDAAM Δ CT and cDAAM^{R-A} originates from their actin nucleation activity, which is intermediate as compared to FH1-FH2 and cDAAM, as revealed by TIRFM measurements (Figure 2, F and 4, C-E).

These observations suggest that DAD contributes to the nucleation promoting activity of cDAAM, but the presence of a functional CT region is absolutely necessary to reconstitute its full nucleation ability. Our data also indicate that the contribution of DAD and CT regions to actin nucleation is non-cooperative.

DAD-CT cannot compensate for a loss-of-function mutation induced defects in the main activities of the FH2 domain

To better understand the role of DAD-CT in DAAM-mediated actin dynamics, the activities of this region were investigated in the presence of a mutant FH2 domain. The α D helix of the knob region of the FH2 domain contains a highly conserved Ile residue, which is essential for FH2-mediated polymerization (Figure 1, B) (13,14,23,35,36). By introducing the I732A mutation, we generated a cDAAM^{I732A} construct that contains the DAD-CT region and an FH1-FH2^{I732A} version which is devoid of the C-terminal domains (Figure 1, A). Steady-state anisotropy measurements revealed that the I732A mutation severely reduces the ability of the FH2 domain to bind both free and profilin:G-actin in the absence of DAD-CT (Figure 5, A, Table 1). However, we observed that cDAAM^{I732A} can bind actin with similar affinity as cDAAM, which indicates that the presence of DAD-CT can compensate for the defects in monomer binding induced by the I732A mutation (Figure 5, A, Table 1).

In pyrenyl polymerization assays we found that FH1-FH2^{I732A} inhibits the overall polymerization of free G-actin, but the assembly rate of profilin:G-actin is not affected in the concentration range that we could test in these experiments (Figure 5, B). The cDAAM^{I732A} mutant is more efficient in reducing the rate of actin assembly from both free G-actin and profilin:G-actin than FH1-FH2^{I732A} (Figure 5, B). Importantly, the inhibition of the bulk polymerization rate by the constructs carrying the mutation is the opposite

as to that was observed for the wild-type FH2 domain (*Figure 2, A and B*). These observations show that the I732A mutation impairs the proper actin assembly activities of DAAM FH2, which is consistent with previous data (36). To better understand the underlying effects, TIRFM experiments were performed. In contrast to the wild type fragments (*Figure 2, E*), filament number was not changed significantly in the presence of either of the two mutant constructs (*Figure 5, C and D*). This finding indicates that despite of being able to interact with actin, the I732A mutation abolishes the nucleation activity of DAAM. Our data also suggest that the presence of the DAD-CT region is able to counteract the actin binding defects of the FH2 domain induced by I732A, however it cannot restore the proper functionality of the mutated FH2 domain in actin nucleation. The analysis of the effects on elongation revealed that FH1-FH2^{I732A} only moderately affects filament growth from free G-actin in the concentration range in which the wild type construct almost completely inhibits elongation ($v_{\text{FH1-FH2}^{\text{I732A}}} = 4.09 \pm 0.4 \text{ subunit} \times \text{s}^{-1}$ ($p = 0.03$), *Figure 5, E*). Even higher amount (20 μM FH1-FH2^{I732A}) resulted only in $\sim 55\%$ inhibition (*Figure 5, E*). In agreement with the pyrenyl polymerization experiments, FH1-FH2^{I732A} does not affect significantly filament growth from profilin:G-actin at low concentrations ($v_{\text{FH1-FH2}^{\text{I732A}}} = 3.53 \pm 0.4 \text{ subunit} \times \text{s}^{-1}$ ($p = 0.30$), *Figure 5, E*), while it moderately inhibits elongation when it is added at higher amounts ($v_{\text{FH1-FH2}^{\text{I732A}}} = 2.09 \pm 0.1 \text{ subunit} \times \text{s}^{-1}$ ($p < 0.0001$), *Figure 5, E*). cDAAM^{I732A} moderately slows filament elongation when it is added at the same amount as cDAAM ($v_{\text{cDAAM}^{\text{I732A}}} = 3.97 \pm 0.3 \text{ subunit} \times \text{s}^{-1}$ ($p = 0.00014$), *Figure 5, E*). At relatively high concentrations (2 μM) it largely slows down the elongation from free G-actin, similarly to the wild-type protein ($v_{\text{cDAAM}^{\text{I732A}}} = 0.15 \pm 0.02 \text{ subunit} \times \text{s}^{-1}$ ($p < 0.0001$), *Figure 5, E*). However, unlike cDAAM that maintains elongation in the presence of profilin (*Figure 2, F*), cDAAM^{I732A} functions oppositely, it hinders profilin:G-actin association to barbed ends in a concentration dependent manner (*Figure 5, E*).

The reduced filament growth rate observed in the presence of cDAAM^{I732A} can result from the altered barbed end rate constants of monomers due to capping-like function and/or a decrease in assembly competent G-actin due to sequestration. To test these

possibilities, the influence of cDAAM^{I732A} on the amount of unassembled actin was measured at steady-state (J(c) plot, *Figure 5, F*). In case of spontaneous actin assembly, the steady-state amount of unassembled free G-actin (critical concentration) is reflected by the breaking point of the J(c) plot (*Figure 5, F*). In the absence of DAAM, the breaking point appeared at $\sim 100 \text{ nM}$, in agreement with the well-established value of the barbed end critical concentration (*Figure 5, F and (1,2)*). The J(c) plot recorded in the presence of cDAAM^{I732A} was parallel to the plot obtained for actin, and the breaking point was not affected significantly (*Figure 5, F*). The complete and permanent blocking of barbed ends by classic barbed end capping proteins, such as CapG would result in a shift of the breaking point up to the value characteristic to the critical concentration of pointed ends ($\sim 600 \text{ nM}$) (1-3,28). The unaffected breaking point indicates that the filament ends are not strongly capped by cDAAM^{I732A} and monomer-polymer exchanges at barbed ends can occur. We made similar observation for other DAAM constructs (21). In case of sequestration, considering the dissociation equilibrium constant of the cDAAM^{I732A}: G-actin interaction (*Table 1*), the critical concentration of actin assembly (*Figure 5, F*) and the concentration of cDAAM^{I732A} used in this experiment (1 μM) one would expect an increase of $\sim 500 \text{ nM}$ in the amount of unassembled actin due to sequestration, which would result in a significant shift in the breaking point to $\sim 600 \text{ nM}$ (28). On the other hand, in case of sequestration the number of actin filaments is expected to be decreased (28), however, in TIRFM experiments we found that cDAAM^{I732A} does not change filament number (*Figure 5, D*), which further supports the lack of sequestering activity. Therefore, these data support that cDAAM^{I732A} does not sequester G-actin, neither blocks completely and permanently monomer-polymer exchange at barbed ends as a classic capping protein. To explain the large but not complete inhibition of elongation by cDAAM^{I732A} one can assume that it slows filament elongation at barbed ends by influencing the rate constants of subunit association and/or dissociation.

Consistently with the importance of the conserved Ile residue in actin interaction (37), the mutation impairs not only the nucleation ability of the FH2 domain but it also diminishes its interaction with the barbed end of the

filaments. The combination of these two effects results in a decreased bulk polymerization rate, detected in the pyrenyl polymerization assays (Figure 5, B). At high concentration, DAAM FH2^{I732A} can maintain some interactions with filament ends similarly to other formins (e.g. mDia1 (13)). Even if such high concentration is physiologically not relevant this approach allowed us to address the effects of DAD-CT on the interaction of FH2 with barbed ends. Our data indicates that besides the key Ile residue, other residues/binding sites in the FH2 domain also contribute to barbed end interactions of DAAM, albeit with much lower affinities. Comparative structural analysis of DAAM FH2 with other formins reveals that residues in the knob region near the Ile732, as well as the lasso/post interface can contribute to actin binding (36). Based on our data the coordination between these different binding sites is crucial for functional barbed end interaction. Importantly, our observations – that the magnitude of the effect of the FH2^{I732A} mutant on filament elongation is more pronounced in the presence of DAD-CT than in the absence of it – suggests that the C-terminal regions are important for filament end interaction.

DAD-CT contributes to the antagonism with Capping Proteins

We found that cDAAM^{I732A} affects filament end dynamics more efficiently than the FH1-FH2^{I732A} (Figure 5, B and E), which suggests that besides nucleation, DAD-CT is also important for filament end interaction. We studied the effect of these domains in the presence of Capping Protein (CP) to further analyze the potential role of DAD-CT in formin:barbed end interaction. CP is well known for its ability to bind to filament barbed ends and block their elongation ((38,39) and Figure 6). Formins were shown to be able to antagonize the effect of CPs at barbed ends to sustain processive growth of actin filaments (40-44). Thus, we investigated whether DAAM can compete with CP for barbed end binding, and if it does so, which regions are important for this activity. Actin assembly was initiated in the absence or presence of different DAAM constructs, then CP in increasing amount was added to the polymerization mixtures after ~ 120 s (Figure 6, A) (40). CP inhibits both the rate and the extent of DAAM-mediated actin assembly for all constructs, however with

different efficiencies (Figure 6). Quantitative analysis revealed that cDAAM is more efficient than FH1-FH2 in protecting filament ends ($IC_{50} = 345.9 \pm 27.60$ nM and $IC_{50} = 47.7 \pm 16.97$ nM, respectively). Mutations in the CT region (cDAAM Δ CT and cDAAM^{R-A}) partially reduce the ability of DAAM to compete with CP ($IC_{50} = 108.6 \pm 19.35$ nM for cDAAM Δ CT, $IC_{50} = 93.7 \pm 19.06$ nM for cDAAM^{R-A} Figure 6). FH1-FH2^{I732A} and cDAAM^{I732A} failed to antagonize the capping effect when added at the same concentration as the wild-type constructs, in agreement with the importance of Ile732 in barbed end interaction (Figure 6, A) (37). Because FH1-FH2^{I732A} and cDAAM^{I732A} do not show any difference in this assay, we conclude that the DAD-CT region is unable to compete for barbed end binding in the absence of a functional FH2 domain (Figure 6, A). Consistent with this, we found that the isolated DAD-CT cannot uncap CP-bound barbed ends (Figure 6, A). These observations corroborate the role of the C-terminal regions of DAAM in filament end interaction. Importantly, our data clearly demonstrate that though the FH2 domain is necessary for uncapping of CP, the DAD-CT regions can contribute to this antagonistic activity by tuning the filament end protection ability of FH2.

The role of DAD-CT in FH2-mediated F-actin interaction

Previously, we showed that both *Drosophila* DAAM FH2 and FH1-FH2 possess actin filament binding and bundling activities (21). This activity is characteristic for the human Daam1 protein, as well (30,45). Based on this, to further extend our studies on the role of DAD-CT in actin interaction, we investigated whether the DAD-CT region can interact with F-actin in high-speed co-sedimentation experiments (Figure 7, A). The amount of DAD-CT sedimented with F-actin increased in a concentration dependent manner, which indicates that DAD-CT can bind to the sides of filamentous actin with a dissociation equilibrium constant of $K_D(\text{DAD-CT}) = 38.9 \pm 3.2$ μ M (Figure 7, A). This suggests less efficient interaction than that of the FH2 domain, which is characterized by K_D in the few micromolar range (21). Removal of the CT significantly reduces the binding strength of the C-terminus ($K_D(\text{DAD}) > 100$ μ M), while the binding of the DAD-CT^{R-A} mutant to F-actin is

negligible (Figure 7, A). To test the functional consequences of side binding, the bundling activity of DAAM was studied in low-speed centrifugation assays. We found that FH1-FH2, as well as cDAAM bundle actin filaments with approximately the same efficiency, and the I723A mutation does not affect the bundling activity of FH2 (Figure 7, B). Our results revealed that the DAD-CT region can also bundle F-actin, independently of the FH2 domain, however, its efficiency is extremely low (Figure 7, B). Neither DAD nor DAD-CT^{R-A} show significant activity at the concentrations we could test in the experiments (Figure 7, B).

Altogether, these results show that the isolated DAD-CT can interact with the sides of actin filaments and also suggest that the FH2 domain of DAAM is the core actin filament side-binding/bundling element, whereas the C-terminal region appears to have a minor contribution to this activity. Consistently with the G-actin interaction, these data further support that the main C-terminal actin-binding element is the CT region.

The actin assembly promoting activity of the FH2 domain of DAAM is regulated through an autoinhibitory interaction

The DRF formin subfamily proteins are regulated by intramolecular autoinhibitory interactions mediated by the N-terminal DID and the C-terminal DAD domains. DAAM possesses the N-, and C-terminal sequence elements characteristic for these regulatory domains (Figure 1, A). Consistently, our previous investigations showed that *Drosophila* DAAM constructs lacking either one of the DID or DAD regions behave as constitutively active forms of the protein in *in vivo* assays (17). To address this issue biochemically, the effects of the recombinant DID fragment on FH1-FH2 (lacking the C-terminal DAD-CT region) and cDAAM (possessing the C-terminal DAD-CT region) mediated actin assembly were investigated in pyrenyl polymerization experiments (Figure 8, A). We found that DID does not affect the spontaneous actin polymerization or the actin assembly promoting activity of FH1-FH2 (Figure 8, A). In contrast, it inhibits the cDAAM mediated actin polymerization in a concentration dependent manner (Figure 8). At saturating DID concentration the polymerization rate was reduced to the value characteristic to

spontaneous actin assembly (Figure 8). For quantitative analysis the relative polymerization rates from the slopes of the pyrenyl traces at half-maximum polymerization were derived, and plotted as the function of DID concentration (Figure 8, B). The fit gave dissociation equilibrium constant of ~ 30 nM, consistent with a tight interaction between the DID and the DAD-CT domains (Eq.1.).

These results support that the interaction of the FH2 domain of DAAM with actin is regulated by intramolecular interactions between the DID and the C-terminal regions (46). Thus, although the DAD-CT domain might be able to bind actin very weakly (Figure 3, A), similar to other DRFs, its major function is likely related to DID binding through which it shades the actin assembly activity of the FH2 domain.

In vivo activity of the CT region

To further assess the functional importance of the actin interacting domains of DAAM, *Drosophila* primary neuronal cultures were used to analyze the *in vivo* effect of FH2 and CT on actin assembly. We took advantage of the fact that growth cone filopodia are primarily actin based structures, and we previously showed that overexpression of a constitutively active form of dDAAM, lacking the DAD domain (Δ DAD-dDAAM), induces a significant increase in filopodia number in primary neurons (17). To exploit this effect, CT-truncated and/or FH2^{I732A}-mutated Δ DAD-dDAAM constructs were expressed in *Drosophila* embryos (Figure 9, A). In accordance with our former results, the pan-neuronal expression of Δ DAD-dDAAM induced a significant increase of axonal filopodia formation as compared to control cells (9.9 ± 0.60 vs. 6.9 ± 0.37) (Figure 9, B and C). In contrast to it, overexpression of Δ DAD-CT-dDAAM induced only a moderate increase in filopodia number (7.8 ± 0.60) (Figure 9, B and C). Thus the constitutively active form of dDAAM was more effective in the presence of the CT region. Contribution of the FH2 domain was analyzed by expression of Δ DAD-dDAAM^{I732A} which was also able to induce a moderate increase in filopodia number (7.9 ± 0.54), comparable to that of Δ DAD-CT-dDAAM (Figure 9, B and C). These results reveal that, despite of the huge differences in their *in vitro* actin assembly activity, the FH2

and CT domains display some requirements during filopodia formation in cultured neurons. In agreement with a partly redundant role, filopodia number of cells expressing Δ DAD-CT-dDAAM^{I732A} was comparable to that of control cells (6.9 ± 0.52) (Figure 9, B and C), thus the lack of CT combined with the I732A mutation abolished completely the actin assembly activity of dDAAM.

To better understand the mechanisms of the actin binding domains of DAAM in filopodia formation, filopodia dynamics was studied in primary neurons expressing Δ DAD-dDAAM or Δ DAD-CT-dDAAM. Filopodia dynamics was followed by co-expression of Actin5C::GFP (Figure 9, D), and as expected, live imaging of 7-9 HIV neurons revealed an extensive movements of filopodia (Supplemental movie 1.) including lateral displacement, collapse, stasis and fusion. To characterize filopodia dynamics we used extension and retraction as readouts (Figure 9, E and F). We found that the overexpression of the truncated DAAM isoforms do not change significantly the extension rate as compared to control neurons, expressing Actin5C::GFP alone (Figure 9, E). However, overexpression of the activated DAAM isoform (Δ DAD-dDAAM) leads to decreased filopodia retraction (Figure 9, F). This finding indicates that activated DAAM is probably involved in filopodia stabilization rather than filopodia elongation, whereas it still remains possible that DAAM is also involved in the initial actin nucleation steps of filopodia formation. The retraction rate upon Δ DAD-CT-dDAAM overexpression is in between the wild type control and that of Δ DAD-dDAAM (Figure 9, F), suggesting that the CT domain may have a minor role in filopodia stabilization.

Thus, the analysis of filopodia number and dynamics in primary neurons revealed that the CT domain of DAAM appears to contribute to filopodia formation *in vivo* and may also affect the dynamic behavior of filopodia.

DISCUSSION

In this work we extended the characterization of the C-terminal elements of formins by analyzing the role of the DAD and CT regions of *Drosophila* DAAM in the regulation of actin dynamics both *in vitro* and *in vivo*. We showed that the DAD-CT tunes the nucleation activity and the proper filament end interactions of the FH1-FH2 domains of DAAM, as well as its ability to antagonize with Capping Protein.

Our data revealed that the DAD-CT region of DAAM can interact with actin *in vitro*, however the binding of isolated DAD is very weak, hence it is the CT region that is largely responsible for actin interaction (Figure 3, A, Table 1). The C-terminal regions of mDia1, INF2 and FMNL3 possess sequence elements characteristic to the N-terminus of the actin-binding WH2 domains (12-14). Comparative sequence analysis of the DAD-CT regions of DAAM indicates some sequence similarity to WH2 domains, as well (Figure 1, C). The DAD region contains the conserved hydrophobic amino acid triplet of LLxxI, which mediates interactions between the N-terminus of WH2 domains and the hydrophobic cleft of actin (34). However, the characteristic LKKT/V motif, which is essential for the actin binding of WH2 domains is completely absent from the C-terminus of DAAM (Figure 1, C). Similar sequence characters can be observed for the DAD of mDia1 and FMNL3, which also bind actin relatively weakly (Figure 1, C, Table 2 and (13,14)). Conversely, INF2 possesses the LKKT/V motif that appears to strengthen its affinity to actin (Table 2 and (12)). The CT of DAAM is a relatively long (~ 40 aa) extension and predicted to be intrinsically disordered. Our mutational analysis indicates that the Arg/Lys residues in the CT region are central to efficient actin binding (Figure 3, A). Basic amino acids in the C-terminus of mDia1, FMNL3 and Capuccino were shown to contribute to their actin affinity (13,14,47). These data suggest that the weak actin binding of the DAD region of formins might be attributed to the hydrophobic WH2-like amino acid triplet, however, other sequence elements that are present in the CT region appear to ensure a significantly stronger interaction.

The DAD-CT region of DAAM has similar affinity for free G-actin and profilin-bound G-actin (Figure 3, A, Table 1). We do not

detect any indication for mutually exclusive binding between profilin and DAD-CT to monomeric actin, while DAD-CT is displaced by WH2 domains possessing relatively long and disordered C-terminal extension (Figure 3, B - C). Profilin interacts with the hydrophobic cleft of subdomain 1 and 3 of actin, which might interfere with the WH2-like binding mode of DAD. The long C-terminal half of WH2 extends along the negative surface patch of the actin molecule towards the pointed face, through interactions mainly controlled by electrostatic forces (32-34). The similar structural features and the competitive binding of DAAM CT and WH2 domains indicate that DAAM CT adapts a binding mode, which is similar to that of the C-terminus of WH2. These considerations further support our conclusion that the main actin binding element of DAD-CT is not the DAD, but the CT region. (Figure 10).

As opposed to mDia1 DAD, the isolated DAD-CT of DAAM cannot nucleate actin filaments on its own, even in its dimeric form maintained by GST (13). However, it markedly contributes to the nucleation efficiency of the FH2 domain, yet a properly functioning FH2 domain is absolutely essential for nucleation (Figure 2, 4, 5). The actin binding strengths of the FH1-FH2 and cDAAM constructs are apparently the same (Figure 2, C, Table 1). From this aspect the FH2 domain of DAAM differs from other formins, which requires the C-terminal regions for efficient G-actin binding (13,22,23). However, the more potent nucleation activity of cDAAM as compared to that of FH1-FH2 indicates that the presence of the DAD-CT region must change the structural and/or kinetic features of FH1-FH2 when in complex with actin. This is supported by our data, which indicates positive cooperativity between the association of FH2 and DAD-CT to actin, as well as structural considerations as discussed below.

Besides their role in stabilizing nucleation intermediers, our results also indicate that the C-terminal regions can contribute to proper filament end interaction and to the processivity of DAAM. This is supported by the fact that cDAAM is more efficient in maintaining filament elongation in the presence of Capping Protein than FH1-FH2, and that the DAD-CT region influences the effect of the I732A mutation on actin elongation (Figure 5, 6). Recent findings on the co-regulation of barbed end dynamics by CPs and

formins (e.g. mDia1, FMNL2 and human Daam1) propose the existence of a ternary, or decision complex at the filament end (40,43). Due to the overlapping binding sites of CPs and formins, the co-existence of these proteins at the filament end results in steric clashes, which can be resolved by the partial dissociation of both CPs and formin from the barbed end, as suggested by structural modelling (43). According to the proposed structural features, the main filament end interacting region of the formin in the ternary complex is the α D helix in the knob region. Disrupting this region, by introducing the mutation in the conserved Ile residue is expected to loosen the formin:barbed end interaction and weaken its integrity in the ternary complex. This may result in a faster dissociation and/or slower association or complete removal of the formin, leaving the ends capped by CPs, consistently with in our experiments (Figure 6). Importantly, our results shed light on the importance of DAD-CT in the anti-capping activity of DAAM. The more efficient anti-capping activity of cDAAM as compared to FH1-FH2 (Figure 6) indicates that the C-terminal regions of DAAM contributes to the stability of the ternary complex, which relies on both DAD and CT. Nevertheless, it seems that the loss-of-function defects in the FH2 domain regarding filament end interaction can be compensated by DAD-CT only in the absence of profilin and CPs. This indicates that in a complex biological context, i.e. filopodial elongation or sarcomeric thin filament lengthening, the FH2 domain of DAAM is essential for properly functioning barbed end interactions.

What is the mechanistic view of the enhancement of FH2-mediated actin assembly by DAD-CT? DAD-CT can stabilize the structure of the FH2 domain making it more efficient nucleator, by which it would contribute indirectly – independent from its own actin binding ability – to the core activity of FH2. A similar mode of action was proposed for the FH1 domain of FMNL3 (48). Alternatively, the binding of isolated DAD-CT to actin suggests that DAD-CT can directly interact with actin in their complexes with FH1-FH2-DAD-CT to promote nucleation. In support of this, structural data predict that each of the DAD-CT regions in the FH1-FH2-DAD-CT dimer can establish contacts with an actin monomer (Figure 10, A). This would result in the stabilization of four actin subunits by DAAM,

which would completely overcome the structural and kinetic barrier of actin assembly imposed by the nucleation phase. Accordingly, the current model proposes that the FH1-FH2-C-terminus forms a tripartite machinery, in which the C-terminus serves as a monomer recruitment motif (13,48). This model implies that the actin monomers captured by DAD-CT would incorporate at pointed ends, which can be interfered by profilin. On the other hand, the model implicates that the affinity of the C-terminal of DAAM and some other formins – which is relatively weak in their isolated forms (Table 2) – must be strengthened in their complexes with FH2. This might occur by FH2-mediated structural changes in the C-terminal regions. Alternatively, the FH2 domain by bringing actin subunits into the close proximity of DAD-CT could increase the apparent affinity of the C-terminus. In this scenario, the low-affinity C-terminal regions of formins may be involved in the stabilization of the FH2:actin complex, while high-affinity C-terminal domains can mediate monomer recruitment (Figure 10, B). Besides nucleation DAD-CT also supports the interaction of FH2 with the filament ends, as well as its anti-capping efficiency. This is manifested possibly through interactions of the DAD-CT with the sub-terminal actin subunits, consistently with the proposed structural model (Figure 10 (48)). In the presence of the C-terminal regions, the stair stepping of formins requires the dissociation and re-association of both FH2 and DAD-CT, which can influence the processive mode of filament end tracking, as suggested by our data and other work (15).

The contribution of the DAD-CT of DAAM in tuning the activities of the FH2 domain is supported by the analysis of its role in neuronal actin dynamics (Figure 9). Our results point to the importance of the CT region in actin interactions, in good agreement with the biochemical data. Interestingly, the *in vivo* data suggest that the contributions of the FH2 and CT regions to filopodia formation are comparable. An explanation might be that the CT region of DAAM is not only important for actin interaction, but it can also associate to other cytoskeletal regulators which can contribute to the functioning of DAAM. As examples, Flightless I was shown to directly interact with the C-terminus of DAAM and modulate its actin assembly activity, also other

formins can bind microtubules which influences their interactions with actin (20,47,49).

In conclusion, the C-terminus of DAAM shares similar properties to other formins with regard to actin binding and tuning the nucleation promoting activity of the FH2 domain, as well as to its processive filament end tracking. We also demonstrate that DAD-CT makes the FH2 domain more efficient in antagonizing with Capping Protein. Our observations suggest that the effects of DAD-CT on the actin nucleation and elongation activities of DAAM are manifested by cooperative structural and/or kinetic stabilization of the interaction of FH2 with actin. Altogether, our work provide support for the idea that the DAD-CT region plays a significant role in modulation of the actin assembling properties of the *Drosophila* member of the DAAM formin.

EXPERIMENTAL PROCEDURES

Protein purifications and modifications

For bacterial protein expression, cDNAs of DAAM subfragments (DID: 115-356 aa, cDAAM: 568-1153 aa, FH1-FH2: 568-1054 aa, cDAAM Δ CT: 568-1116, DAD-CT: 1083-1153 aa, DAD: 1083-1119 aa) and their mutated versions (FH1-FH2^{I732A}, cDAAM^{I732A}, cDAAM^{R-A}, DAD-CT^{R-A}) (Figure 1, A) were inserted into pGEX-2T vector (Amersham Biosciences). Constructs were expressed as Glutathion S-Transferase (GST) fusion proteins in *Escherichia coli* BL21(DE3)pLysS strain (Novagen). Transformed bacteria were grown at 37°C in Luria Broth powder microbial growth medium (Sigma-Aldrich). Protein expression was induced by addition of 1 mM isopropyl β -D-1-thiogalactopyranoside (IPTG) at OD₆₀₀ ~ 0.6 - 0.8. After overnight expression at 20°C the bacterial extracts were collected by centrifugation (10.000 g, 15 min, 4°C) and stored at - 80°C until use. For protein purification the bacterial pellet was lysed by sonication in Lysis buffer (50 mM Tris-HCl pH7.6, 5 mM DTT, 50 mM NaCl, 5 mM EDTA, 1 % sucrose, 10 % glycerol supplemented with 1 mM PMSF, 5 mM MgCl₂, 0.1 mg/ml DNase and Protease Inhibitor Cocktail (Sigma-Aldrich P8465)). The cell lysate was ultracentrifuged (110.000 g, 1 h, 4°C). The supernatant was slowly loaded onto GSH resin (Amersham Biosciences) in column. For Formin homology (FH) domain containing constructs, the column was sequentially washed with Lysis, Wash1 (50 mM Tris-HCl pH 7.6, 5 mM DTT, 400 mM NaCl, 10 % (v/v) glycerol, 1% (w/v) sucrose), ATP (50 mM Tris-HCl pH 7.6, 5 mM DTT, 100 mM KCl, 10 mM MgCl₂, 0.25 mM ATP, 5 % (v/v) glycerol, 1 % (w/v) sucrose) and Wash2 (50 mM Tris-HCl pH 7.6, 5 mM DTT, 50 mM NaCl, 5 mM MgCl₂, 10 mM KCl, 5 % (v/v) glycerol, 1 % (w/v) sucrose) buffer. For the C-terminal constructs, lacking the FH domains, the column was washed with Lysis and Wash2 buffer. The proteins were eluted by Wash2 buffer supplemented with 50 mM Glutathion Reduced (Sigma G4251), concentrated (Vivaspin 30.000 Da cut-off) and loaded onto PD10 column (GE Healthcare 17-0851-01) for buffer exchange to Storing buffer (50 mM Hepes pH 7.6, 5 mM DTT, 50 mM NaCl, 5 % (v/v) glycerol, 1 % (w/v) sucrose). Before flash freeze in liquid nitrogen the constructs were clarified by ultracentrifugation (300.000 g, 30

min, 4°C) and stored at - 80°C until use. Control experiments showed that a freeze/thaw cycle does not affect the functionality of the constructs (*data not shown*). Typically, 5 – 6 g bacterial pellet yielded 1 – 1.2 mg/ml protein. The protein concentrations were measured spectrophotometrically using the extinction coefficients at 280 nm and molecular weights derived from the amino acid sequence (Expasy ProtParam tool <http://web.expasy.org/protparam/>). Actin was purified from rabbit skeletal muscle, gel filtered on Superdex 200 and stored in G buffer (4 mM Tris-HCl pH 7.8, 0.1 mM CaCl₂, 0.2 mM ATP, 0.005 % NaN₃, 0.5 mM beta-mercaptoethanol) according to standard protocols (50,51). Actin was labelled at Lys³²⁸ by Alexa Fluor[®] 488 carboxylic acid succinimidyl ester (Alexa488NHS, Invitrogen) or at Cys³⁷⁴ by N-(1-Pyrene)Iodoacetamide (pyrene, Thermo Fischer Scientific) according to standard protocols (21,28,52). Mouse profilin 1 was purified as described previously and labelled with Alexa Fluor[®] C₅ 568 maleimide (Alexa568C, Invitrogen) (28,53). Heterodimeric mouse $\alpha\beta$ Capping Protein (CP) and the WH2 domain-containing construct of SALS (SALS-WH2) were purified as described previously (28,54).

General experimental considerations

Samples at each concentration were prepared individually for experiments. All measurements were performed at 20°C. The sum of the volume of the proteins and the volume of their storing buffer was constant in the samples and represented maximum 50 % of the total volume of the sample. The concentrations given in the text are final concentrations. In all experiments Mg²⁺-ATP- actin was used. The actin monomer bound Ca²⁺ was replaced by Mg²⁺ by adding 200 μ M EGTA and 50 μ M MgCl₂ and incubating the samples for 5 – 10 min at room temperature.

Pyrenyl polymerization assays

The polymerization kinetics of Mg²⁺-ATP-G-actin was measured using pyrene-actin. The actin concentration was 2 μ M that contained 5 % or 2 % pyrenyl labelled actin in case of free G-actin and profilin-G-actin, respectively. In profilin containing samples the profilin concentration was 6 μ M. The polymerization was initiated by the addition of 1 mM MgCl₂ and 50 mM KCl in the absence and presence of

different proteins (for exact sample composition and concentrations see *Figure Legends*). The measurements were performed using a Safas Xenius FLX spectrofluorimeter ($\lambda_{\text{ex}} = 365 \text{ nm}$, $\lambda_{\text{em}} = 407 \text{ nm}$). To quantitatively analyze the effect of DAAM on actin assembly, the polymerization rates were determined from the slope of the pyrenyl traces at half-maximum polymerization. The relative polymerization rates were derived as the ratio of the polymerization rate measured in the presence of different amounts of DAAM and the polymerization rate of spontaneous actin assembly (*Figure 2, B*, *Figure 4, B*). For the quantitative analysis of the effect of DID on cDAAM mediated actin assembly (*Figure 8, B*), the relative polymerization rate was derived as the ratio of the polymerization rate measured in the presence of cDAAM and different amounts of DID and the polymerization rate measured in the presence of cDAAM and in the absence of DID. The DID concentration dependence of the relative polymerization rate (v_{relative}) was fit by the following equation:

$$v_{\text{relative}} = \left(1 - \frac{[\text{cDAAM:DID}]}{[\text{cDAAM}_0]}\right) * v_0 + v_{\text{min}} \quad (\text{Eq. 1.})$$

where v_0 and v_{min} is the relative polymerization rate in the absence and presence of saturating amount of DID, respectively, $[\text{cDAAM}_0]$ is the total cDAAM concentration and $[\text{cDAAM:DID}]$ is the concentration of the cDAAM:DID complex, which was derived from the quadratic binding equation.

The antagonistic effect of DAAM with CP was investigated as described (40). Briefly, profilin:G-actin assembly was initiated in the absence of CP and in the presence of different DAAM constructs (200 nM), then CP at different concentrations was added after ~ 120 s to the protein mixtures. Polymerization rates (v) were derived over 40 s just before and 40 s after the addition of CP. The rates were plotted as the function of $[\text{CP}]$, data were fit by the following equation:

$$v = v_0 - \left(\frac{v_0 - v_{\text{min}}}{1 + \frac{[\text{CP}]}{IC_{50}}}\right) \quad (\text{Eq. 2.})$$

where v_0 and v_{min} is the polymerization rate in the absence and presence of saturating amount of CP, respectively, $[\text{CP}_0]$ is the total CP

concentration and IC_{50} is the CP concentration required for 50 % inhibition.

Steady-state fluorescence experiments

Steady-state anisotropy: To study the DAAM:G-actin interaction the steady-state anisotropy of AlexaFluor488 succinimidyl ester labelled Mg^{2+} -ATP-G-actin (Alexa488NHS-G-actin) was measured. Alexa488NHS-G-actin (0.2 μM) was incubated with LatrunculinA (LatA, 4 μM) for 20 min. Then DAAM constructs were added at different concentrations and the samples were further incubated for 1 h at 20°C. In measurements when profilin was present, profilin (0.8 μM) was added to actin after the incubation with LatA, and the samples were further incubated for 30 min at 20°C, prior to the addition of DAAM constructs. Note, that due to the presence of LatA that prevents actin polymerization, the increase in steady-state anisotropy could not result from filament formation, it solely reflects the binding of DAAM constructs to actin. To study the salt dependence of the DAD-CT:G-actin interaction, the ionic strength was set by adding NaCl to the samples (for exact concentrations see *Figure 3, A*). Anisotropy measurements were performed using a Horiba Jobin Yvon spectrofluorometer ($\lambda_{\text{ex}} = 488 \text{ nm}$, $\lambda_{\text{em}} = 516 \text{ nm}$, slits: 5 nm). To study the interaction of DAD-CT with profilin:G-actin, steady-state anisotropy measurements were performed using AlexaFluor568C₅ maleimide labelled profilin (Alexa568C-profilin). In these experiments Alexa568C-profilin (2 μM) was added to Mg^{2+} -ATP-G-actin (4 μM) bound to LatA (8 μM), after 30 min incubation at 20°C the samples were supplemented with DAD-CT at different concentrations and further incubated for 1 h at 20°C. The measurements were performed using a Horiba Jobin Yvon spectrofluorometer ($\lambda_{\text{ex}} = 578 \text{ nm}$, $\lambda_{\text{em}} = 600 \text{ nm}$, slits: 5 nm). For quantitative analysis the DAAM concentration dependence of the steady-state anisotropy (r) measured either in the absence or presence of profilin was fit by the following equation:

$$\frac{r - r_A}{r_{AD} - r_A} = \frac{A_0 + D_0 + K_d - \sqrt{(A_0 + D_0 + K_d)^2 - 4A_0D_0}}{2D_0} \quad (\text{Eq. 3.})$$

where A_0 and D_0 are the total G-actin and DAAM concentration, respectively, r_A is the

steady-state anisotropy of Alexa488NHS-G-actin/Alexa568C-profilin, r_{AD} is the steady-state anisotropy of Alexa488NHS-G-actin/Alexa568C-profilin at saturating amount of DAAM, K_d is the dissociation equilibrium constant of the G-actin : DAAM complex.

J(c) plot measurements: To investigate the effect of cDAAM^{I732A} on the amount of unassembled actin at steady-state critical concentration measurements were performed as described (21,28). The cDAAM^{I732A} concentration was 1 μ M. For the analysis J(c) plots were generated ([actin] dependence of the fluorescence emission).

Total internal reflection fluorescence microscopy

The effects of DAAM constructs on the assembly of individual actin filaments were followed by total internal reflection fluorescence microscopy (TIRFM) as described previously (21,28). Glass flow cells were incubated with 1 volume of N-ethylmaleimide myosin for 1 min, then washed with 2 volumes of myosin buffer (4 mM Tris-HCl pH 7.8, 1 mM DTT, 0.2 mM ATP, 0.1 mM CaCl₂, 500 mM KCl, 1 mM MgCl₂, 0.2 mM EGTA) and 2 volumes of 0.1 % (w/v) BSA. Finally, flow cells were equilibrated with 2 volumes of TIRFM buffer (0.5 % (w/v) methylcellulose, 0.5 % (w/v) BSA, 50 mM 1,4-diazabicyclo-[2,2,2]octane (DABCO), 100 mM DTT in buffer F* (4 mM Tris-HCl, pH 7.8, 1 mM DTT, 0.2 mM ATP, 0.1 mM CaCl₂, 50 mM KCl, 1 mM MgCl₂, 0.2 mM EGTA)) before adding the protein mixture (for exact protein composition and concentration see *Figure Legends*). To follow the assembly of free G-actin or profilin:G-actin in the absence and presence of DAAM a mixture of G-actin (0.5 μ M, containing 10 % Alexa488NHS-G-actin) and different DAAM constructs in TIRFM buffer was injected into the flow cell. In profilin containing samples the profilin concentration was 2 μ M. Images were captured every 10 s with an Olympus IX81 microscope equipped with a laser-based (491 nm) TIRFM module using an APON TIRF 60x NA1.45 oil immersion objective and a CCD camera (Hamamatsu). Images were background corrected before analysis. For the analysis of the number of filaments images were captured 15 min after the initiation of actin polymerization. Filament number was derived from a 66 \times 66 μ m region of the images by using Fiji. Time-lapse

images were analyzed by either the MultipleKymograph plugin of Fiji or by manually tracking filament growth to derive the elongation rate of actin filaments. Filament length was converted to subunits using 370 subunits per micron filament (55). The elongation rate of actin filaments (v) was related to the critical concentration of actin assembly ($cc \sim 0.1 \mu$ M, (1,2)), to the association rate constant of actin monomers to filament barbed ends (k_+) and to the total actin concentration ($[G_0]$) by the following equation:

$$v = k_+([G_0] - cc) \quad (Eq. 4.)$$

Actin filament binding and bundling assays

To investigate the interaction of DAD-CT with actin filaments high-speed centrifugation experiments were performed and analyzed as described (21,28). In control experiments we found that the C-terminal constructs appear in the pellet in the absence of F-actin, however, at significantly lower amount than in the presence of actin filaments. Therefore, for the quantitative analysis the amount of self-pelleting DAAM protein was subtracted from the amount of DAAM sedimented in the presence of actin. The final F-actin concentration was 2.5 μ M. To study the bundling activity of DAAM constructs, Mg²⁺-ATP-G-actin (10 μ M) was polymerized for 2 h at 20°C by adding 1 mM MgCl₂ and 50 mM KCl. Actin filaments were diluted to 1 μ M in F buffer (G buffer supplemented with 1 mM MgCl₂ and 50 mM KCl), in the absence and presence of different DAAM constructs and further incubated for 1 h at 20°C. Samples were centrifuged (14.000 g, 5 min, 20°C), then the supernatants were carefully removed and processed for SDS-PAGE analysis. The protein content of the supernatants was derived from Coomassie blue stained gels (Syngene bioimaging system). Quantification of Coomassie blue intensities were performed within the linear range of exposure identified by a calibration curve. The intensity values were corrected for the molecular weight of each protein. The relative amount of F-actin in the supernatant was derived as the ratio of the amount F-actin in the absence of any other protein to the amount F-actin in the presence of different proteins.

Genetics

The following fly stocks were used for *in vivo* overexpression: *ElavGAL4^{C155}* or *w;elavGAL4* were crossed to *w;UASADAD-DAAM*, *w;UASADAD-CT-DAAM*, *w;UASADAD-DAAM I732A*, *w;UASADAD-CT-DAAM I732A*, *w;UAS-ADAD-DAAM;UAS-Actin5C::GFP*, *w;UAS-ADAD-CT-DAAM;UAS-Actin5C::GFP*.

Primary cell cultures

Drosophila primary neuronal cells were obtained from stage 11 embryos following a protocol published by Sanchez-Soriano N et al. (56) with some modifications. In brief, whole embryos were squashed in Schneider's *Drosophila* (Sigma-Aldrich) medium supplemented with 20 % heat-inactivated fetal bovine serum (Sigma-Aldrich), 2 µg/mL insulin (Sigma-Aldrich) and penicillin-streptomycin solution (Lonza). Embryonic homogenates were spun at 380g for 4 min at room temperature. Pellets were re-suspended in complete medium and the cells were grown at 27°C on glass coverslips or in glass-bottom Petri-dishes for live imaging (MaTek). For filopodia number analysis, primary neurons were fixed 6h after plating in 4 % formaldehyde for 30 min at room temperature. Cells were washed with PBST (PBS + 0.1 % TritonX-100), then blocked in 5 % goat-serum diluted in PBST for 10 min. Cells were incubated with primary antibodies for overnight at 4°C (mouse anti-tubulin 1:1000, (Sigma-Aldrich) rat anti-actin 1:200 (Babraham Bioscience), rabbit anti-HRP 1:200 (Jackson ImmunoResearch), diluted in blocking solution). After overnight incubation cells were washed in PBST, then incubated with secondary antibodies for 1h at room temperature (anti-mouse-Alexa405 1:600, anti-rat-Alexa488 1:600, anti-rabbit-Alexa647 1:600 (Life Technologies) diluted in blocking solution). Cells were washed in PBST and then coverslips were mounted on microscope slides

in 70 % glycerol. Confocal images were collected with an Olympus FV1000 LSM microscope and edited with ImageJ software.

Live imaging and filopodia dynamics

Filopodia dynamics measurements were performed on 7-9 HIV neurons in culture using LSM 880 confocal microscope (Zeiss) equipped with a 40X, 1.4 NA oil-immersion lens. To follow actin dynamics, we overexpressed Actin5C::GFP in the cell culture, using a pan-neuronal driver (*Elav-Gal4*). Imaging of the neurons was performed in glass bottom petri dishes (MatTek corporation) in growth media. Fluorescence was excited using the 488 nm line of the argon laser and recorded at a bandwidth of 500–550 nm at an acquisition rate of 1.27 Hz. Filopodia with recognizable extension and retraction phases were selected for further analysis. To measure the extension and retraction speed, kymographs were built using KymoResliceWide, a Fiji plugin dedicated to generate kymographs with improved contrast. Steepness was measured manually and converted to extension and retraction velocities in Excel (Microsoft).

Statistics

The data presented were derived from at least two independent experiments. Values are displayed as mean ± standard deviation. The number of independent experiments are given in *Figure Legends*. The significance was calculated using two-sample T-tests or Z-test considering the number of data and the variance (Excel, Microsoft). The data were analyzed by Kruskal Wallis or one-way analysis of variance (ANOVA) considering the distribution of the data. By convention $p \geq 0.05$ was considered as statistically not significant, $*p < 0.05$, $**p < 0.01$, $***p < 0.001$, $****p < 0.0001$. The significance levels are given in the text and on the corresponding figure.

ACKNOWLEDGEMENTS

This research was supported by grants from the Hungarian Science Foundation (OTKA) Grants K109689 (to BB) and K109330 (to JM), the Hungarian Brain Research Program (KTIA_NAP_13-2-2014-0007) (to JM), the National Research, Development and Innovation Office (GINOP-2.3.2-15-2016-00001) (to JM), the National Innovation Office “Baross Gábor” Program (REG-DD-09-1-2009-0009 Tírfm 09 (to BB)). This research was supported by the European Union and the State of Hungary, co-financed by the European Social Fund in the framework of TÁMOP 4.2.4.A/2-11-1-2012-0001 ‘National Excellence Program’ and the New National Excellence Program of the Ministry of Human Capacities (to BB and TH). Supported by the ÚNKP-16-4 New National Excellence Program of the Ministry of Human Capacities (to BB).

CONFLICT OF INTEREST

The authors declare that they have no conflicts of interest with the contents of this article.

AUTHORS CONTRIBUTIONS

ATV performed the experiments, analyzed the data, prepared the digital images, IF performed the experiments, analyzed the data, prepared the digital images, SzSz performed the experiments, analyzed the data, prepared the digital images, EM performed the experiments, RG performed the experiments, analyzed the data, MÁT, TH, RP performed the experiments, analyzed the data, GCST read the paper for intellectual content, JM conceived and design the experiments, analyzed the data, drafted the article, BB conceived and design the experiments, performed the experiments, analyzed the data, drafted the article, prepared the digital images.

REFERENCES

1. Bugyi, B., and Carlier, M. F. (2010) Control of actin filament treadmilling in cell motility. *Annu Rev Biophys* **39**, 449-470
2. Pollard, T. D. (2007) Regulation of actin filament assembly by Arp2/3 complex and formins. *Annu Rev Biophys Biomol Struct* **36**, 451-477
3. Pollard, T. D. (2016) Actin and Actin-Binding Proteins. *Cold Spring Harb Perspect Biol* **8**
4. Schonichen, A., and Geyer, M. (2010) Fifteen formins for an actin filament: a molecular view on the regulation of human formins. *Biochim Biophys Acta* **1803**, 152-163
5. Goode, B. L., and Eck, M. J. (2007) Mechanism and function of formins in the control of actin assembly. *Annu Rev Biochem* **76**, 593-627
6. Breitsprecher, D., and Goode, B. L. (2013) Formins at a glance. *J Cell Sci* **126**, 1-7
7. Alberts, A. S. (2001) Identification of a carboxyl-terminal diaphanous-related formin homology protein autoregulatory domain. *J Biol Chem* **276**, 2824-2830
8. Li, F., and Higgs, H. N. (2003) The mouse Formin mDia1 is a potent actin nucleation factor regulated by autoinhibition. *Curr Biol* **13**, 1335-1340
9. Otomo, T., Otomo, C., Tomchick, D. R., Machius, M., and Rosen, M. K. (2005) Structural basis of Rho GTPase-mediated activation of the formin mDia1. *Mol Cell* **18**, 273-281
10. Otomo, T., Tomchick, D. R., Otomo, C., Machius, M., and Rosen, M. K. (2010) Crystal structure of the Formin mDia1 in autoinhibited conformation. *PLoS One* **5**
11. Rose, R., Weyand, M., Lammers, M., Ishizaki, T., Ahmadian, M. R., and Wittinghofer, A. (2005) Structural and mechanistic insights into the interaction between Rho and mammalian Dia. *Nature* **435**, 513-518
12. Chhabra, E. S., and Higgs, H. N. (2006) INF2 Is a WASP homology 2 motif-containing formin that severs actin filaments and accelerates both polymerization and depolymerization. *J Biol Chem* **281**, 26754-26767
13. Gould, C. J., Maiti, S., Michelot, A., Graziano, B. R., Blanchoin, L., and Goode, B. L. (2011) The formin DAD domain plays dual roles in autoinhibition and actin nucleation. *Curr Biol* **21**, 384-390
14. Heimsath, E. G., Jr., and Higgs, H. N. (2012) The C terminus of formin FMNL3 accelerates actin polymerization and contains a WH2 domain-like sequence that binds both monomers and filament barbed ends. *J Biol Chem* **287**, 3087-3098
15. Vizcarra, C. L., Bor, B., and Quinlan, M. E. (2014) The role of formin tails in actin nucleation, processive elongation, and filament bundling. *J Biol Chem* **289**, 30602-30613
16. Matussek, T., Djiane, A., Jankovics, F., Brunner, D., Mlodzik, M., and Mihaly, J. (2006) The Drosophila formin DAAM regulates the tracheal cuticle pattern through organizing the actin cytoskeleton. *Development* **133**, 957-966
17. Matussek, T., Gombos, R., Szecsenyi, A., Sanchez-Soriano, N., Czibula, A., Pataki, C., Gedai, A., Prokop, A., Rasko, I., and Mihaly, J. (2008) Formin proteins of the DAAM subfamily play a role during axon growth. *J Neurosci* **28**, 13310-13319
18. Molnar, I., Migh, E., Szikora, S., Kalmar, T., Vegh, A. G., Deak, F., Barko, S., Bugyi, B., Orfanos, Z., Kovacs, J., Juhasz, G., Varo, G., Nyitrai, M., Sparrow, J., and Mihaly, J. (2014) DAAM is required for thin filament formation and Sarcomerogenesis during muscle development in Drosophila. *PLoS Genet* **10**, e1004166
19. Bao, B., Zhang, L., Hu, H., Yin, S., and Liang, Z. (2012) Deletion of a single-copy DAAM1 gene in congenital heart defect: a case report. *BMC Med Genet* **13**, 63
20. Vogler, G., Liu, J., Iafe, T. W., Migh, E., Mihaly, J., and Bodmer, R. (2014) Cdc42 and formin activity control non-muscle myosin dynamics during Drosophila heart morphogenesis. *J Cell Biol* **206**, 909-922

21. Barko, S., Bugyi, B., Carlier, M. F., Gombos, R., Matusek, T., Mihaly, J., and Nyitrai, M. (2010) Characterization of the biochemical properties and biological function of the formin homology domains of Drosophila DAAM. *J Biol Chem* **285**, 13154-13169
22. Pring, M., Evangelista, M., Boone, C., Yang, C., and Zigmond, S. H. (2003) Mechanism of formin-induced nucleation of actin filaments. *Biochemistry* **42**, 486-496
23. Xu, Y., Moseley, J. B., Sagot, I., Poy, F., Pellman, D., Goode, B. L., and Eck, M. J. (2004) Crystal structures of a Formin Homology-2 domain reveal a tethered dimer architecture. *Cell* **116**, 711-723
24. Coue, M., Brenner, S. L., Spector, I., and Korn, E. D. (1987) Inhibition of actin polymerization by latrunculin A. *FEBS letters* **213**, 316-318
25. Bosch, M., Le, K. H., Bugyi, B., Correia, J. J., Renault, L., and Carlier, M. F. (2007) Analysis of the function of Spire in actin assembly and its synergy with formin and profilin. *Mol Cell* **28**, 555-568
26. Pollard, T. D., and Cooper, J. A. (1984) Quantitative analysis of the effect of Acanthamoeba profilin on actin filament nucleation and elongation. *Biochemistry* **23**, 6631-6641
27. Renault, L., Bugyi, B., and Carlier, M. F. (2008) Spire and Cordon-bleu: multifunctional regulators of actin dynamics. *Trends Cell Biol* **18**, 494-504
28. Toth, M. A., Majoros, A. K., Vig, A. T., Migh, E., Nyitrai, M., Mihaly, J., and Bugyi, B. (2016) Biochemical Activities of the Wiskott-Aldrich Syndrome Homology Region 2 Domains of Sarcomere Length Short (SALS) Protein. *J Biol Chem* **291**, 667-680
29. Pollard, T. D. (1986) Rate constants for the reactions of ATP- and ADP-actin with the ends of actin filaments. *J Cell Biol* **103**, 2747-2754
30. Jaiswal, R., Breitsprecher, D., Collins, A., Correa, I. R., Jr., Xu, M. Q., and Goode, B. L. (2013) The formin Daam1 and fascin directly collaborate to promote filopodia formation. *Curr Biol* **23**, 1373-1379
31. Schutt, C. E., Myslik, J. C., Rozycki, M. D., Goonesekere, N. C., and Lindberg, U. (1993) The structure of crystalline profilin-beta-actin. *Nature* **365**, 810-816
32. Didry, D., Cantrelle, F. X., Husson, C., Roblin, P., Moorthy, A. M., Perez, J., Le Clainche, C., Hertzog, M., Guittet, E., Carlier, M. F., van Heijenoort, C., and Renault, L. (2012) How a single residue in individual beta-thymosin/WH2 domains controls their functions in actin assembly. *Embo J* **31**, 1000-1013
33. Dominguez, R. (2016) The WH2 Domain and Actin Nucleation: Necessary but Insufficient. *Trends Biochem Sci* **41**, 478-490
34. Renault, L., Deville, C., and van Heijenoort, C. (2013) Structural features and interfacial properties of WH2, beta-thymosin domains and other intrinsically disordered domains in the regulation of actin cytoskeleton dynamics. *Cytoskeleton (Hoboken)* **70**, 686-705
35. Shimada, A., Nyitrai, M., Vetter, I. R., Kuhlmann, D., Bugyi, B., Narumiya, S., Geeves, M. A., and Wittinghofer, A. (2004) The core FH2 domain of diaphanous-related formins is an elongated actin binding protein that inhibits polymerization. *Mol Cell* **13**, 511-522
36. Lu, J., Meng, W., Poy, F., Maiti, S., Goode, B. L., and Eck, M. J. (2007) Structure of the FH2 domain of Daam1: implications for formin regulation of actin assembly. *J Mol Biol* **369**, 1258-1269
37. Otomo, T., Tomchick, D. R., Otomo, C., Panchal, S. C., Machius, M., and Rosen, M. K. (2005) Structural basis of actin filament nucleation and processive capping by a formin homology 2 domain. *Nature* **433**, 488-494
38. Isenberg, G., Aebi, U., and Pollard, T. D. (1980) An actin-binding protein from Acanthamoeba regulates actin filament polymerization and interactions. *Nature* **288**, 455-459
39. Wear, M. A., Yamashita, A., Kim, K., Maeda, Y., and Cooper, J. A. (2003) How capping protein binds the barbed end of the actin filament. *Curr Biol* **13**, 1531-1537
40. Bombardier, J. P., Eskin, J. A., Jaiswal, R., Correa, I. R., Jr., Xu, M. Q., Goode, B. L., and Gelles, J. (2015) Single-molecule visualization of a formin-capping protein 'decision complex' at the actin filament barbed end. *Nat Commun* **6**, 8707

41. Ramabhadran, V., Gurel, P. S., and Higgs, H. N. (2012) Mutations to the formin homology 2 domain of INF2 protein have unexpected effects on actin polymerization and severing. *J Biol Chem* **287**, 34234-34245
42. Romero, S., Le Clainche, C., Didry, D., Egile, C., Pantaloni, D., and Carlier, M. F. (2004) Formin is a processive motor that requires profilin to accelerate actin assembly and associated ATP hydrolysis. *Cell* **119**, 419-429
43. Shekhar, S., Kerleau, M., Kuhn, S., Pernier, J., Romet-Lemonne, G., Jegou, A., and Carlier, M. F. (2015) Formin and capping protein together embrace the actin filament in a menage a trois. *Nat Commun* **6**, 8730
44. Harris, E. S., Li, F., and Higgs, H. N. (2004) The mouse formin, FRLalpha, slows actin filament barbed end elongation, competes with capping protein, accelerates polymerization from monomers, and severs filaments. *J Biol Chem* **279**, 20076-20087
45. Higashi, T., Ikeda, T., Shirakawa, R., Kondo, H., Kawato, M., Horiguchi, M., Okuda, T., Okawa, K., Fukai, S., Nureki, O., Kita, T., and Horiuchi, H. (2008) Biochemical characterization of the Rho GTPase-regulated actin assembly by diaphanous-related formins, mDia1 and Daam1, in platelets. *J Biol Chem* **283**, 8746-8755
46. Liu, W., Sato, A., Khadka, D., Bharti, R., Diaz, H., Runnels, L. W., and Habas, R. (2008) Mechanism of activation of the Formin protein Daam1. *Proc Natl Acad Sci U S A* **105**, 210-215
47. Roth-Johnson, E. A., Vizcarra, C. L., Bois, J. S., and Quinlan, M. E. (2014) Interaction between microtubules and the Drosophila formin Cappuccino and its effect on actin assembly. *J Biol Chem* **289**, 4395-4404
48. Thompson, M. E., Heimsath, E. G., Gauvin, T. J., Higgs, H. N., and Kull, F. J. (2013) FMNL3 FH2-actin structure gives insight into formin-mediated actin nucleation and elongation. *Nat Struct Mol Biol* **20**, 111-118
49. Higashi, T., Ikeda, T., Murakami, T., Shirakawa, R., Kawato, M., Okawa, K., Furuse, M., Kimura, T., Kita, T., and Horiuchi, H. (2010) Flightless-I (Fli-I) regulates the actin assembly activity of diaphanous-related formins (DRFs) Daam1 and mDia1 in cooperation with active Rho GTPase. *J Biol Chem* **285**, 16231-16238
50. Spudich, J. A., and Watt, S. (1971) The regulation of rabbit skeletal muscle contraction. I. Biochemical studies of the interaction of the tropomyosin-troponin complex with actin and the proteolytic fragments of myosin. *J Biol Chem* **246**, 4866-4871
51. Bugyi, B., Didry, D., and Carlier, M. F. (2010) How tropomyosin regulates lamellipodial actin-based motility: a combined biochemical and reconstituted motility approach. *Embo J*
52. Bugyi, B., Papp, G., Hild, G., Lorinczy, D., Nevalainen, E. M., Lappalainen, P., Somogyi, B., and Nyitrai, M. (2006) Formins regulate actin filament flexibility through long range allosteric interactions. *J Biol Chem* **281**, 10727-10736
53. Perelroizen, I., Marchand, J. B., Blanchoin, L., Didry, D., and Carlier, M. F. (1994) Interaction of profilin with G-actin and poly(L-proline). *Biochemistry* **33**, 8472-8478
54. Nag, S., Ma, Q., Wang, H., Chumnarnsilpa, S., Lee, W. L., Larsson, M., Kannan, B., Hernandez-Valladares, M., Burtnick, L. D., and Robinson, R. C. (2009) Ca²⁺ binding by domain 2 plays a critical role in the activation and stabilization of gelsolin. *Proc Natl Acad Sci U S A* **106**, 13713-13718
55. Hanson, J., and Lowy, J. (1963) The structure of F-actin and of actin filaments isolated from muscle. *Journal of Molecular Biology* **6**, 46-60
56. Sanchez-Soriano, N., Travis, M., Dajas-Bailador, F., Goncalves-Pimentel, C., Whitmarsh, A. J., and Prokop, A. (2009) Mouse ACF7 and drosophila short stop modulate filopodia formation and microtubule organisation during neuronal growth. *J Cell Sci* **122**, 2534-2542
57. Liu, W., Xie, Y., Ma, J., Luo, X., Nie, P., Zuo, Z., Lahrmann, U., Zhao, Q., Zheng, Y., Zhao, Y., Xue, Y., and Ren, J. (2015) IBS: an illustrator for the presentation and visualization of biological sequences. *Bioinformatics* **31**, 3359-3361
58. Oda, T., Iwasa, M., Aihara, T., Maeda, Y., and Narita, A. (2009) The nature of the globular- to fibrous-actin transition. *Nature* **457**, 441-445

59. Yamashita, M., Higashi, T., Suetsugu, S., Sato, Y., Ikeda, T., Shirakawa, R., Kita, T., Takenawa, T., Horiuchi, H., Fukai, S., and Nureki, O. (2007) Crystal structure of human DAAM1 formin homology 2 domain. *Genes Cells* **12**, 1255-1265
60. Lammers, M., Rose, R., Scrima, A., and Wittinghofer, A. (2005) The regulation of mDia1 by autoinhibition and its release by Rho*GTP. *Embo J* **24**, 4176-4187
61. Chereau, D., Kerff, F., Graceffa, P., Grabarek, Z., Langsetmo, K., and Dominguez, R. (2005) Actin-bound structures of Wiskott-Aldrich syndrome protein (WASP)-homology domain 2 and the implications for filament assembly. *Proc Natl Acad Sci U S A* **102**, 16644-16649

TABLES

Table 1. Dissociation equilibrium constants (K_D) of the DAAM:actin interactions.

construct	$K_D \pm SD^1$ (μM)		
	G-actin	profilin:G-actin ^{FL}	profilin ^{FL} :G-actin
FH1-FH2	0.061 ± 0.02	0.056 ± 0.02	nd
cDAAM	0.052 ± 0.02	0.068 ± 0.02	nd
DAD-CT	3.87 ± 0.19	3.54 ± 0.22	4.71 ± 0.53
	9.71 ± 0.51 (17 mM NaCl)	nd	nd
	44.43 ± 2.85 (50 mM NaCl)	nd	nd
DAD	> 100	> 100	nd
DAD-CT^{R-A}	> 500	> 500	nd
FH1-FH2^{I732A}	6.78 ± 0.96	4.41 ± 0.76	nd
cDAAM^{I732A}	0.064 ± 0.02	0.093 ± 0.04	nd

¹The values were derived in the presence of 5 mM NaCl, unless it is indicated otherwise.

FL: fluorescently labelled protein

Table 2. Properties of the C-terminal elements of different formins.

formin	main actin interacting module ¹	G-actin binding affinity	nucleation activity in isolated form/ contribution to FH2-mediated nucleation	affecting G-actin interaction	other effects in actin dynamics
<i>Mm</i> INF2 (12)	WH2-like/DAD sequence	$K_D \sim 0.06 \mu M$ 50 mM KCl	no/yes	profilin interferes	monomer sequestration, filament severing
<i>Mm</i> FMNL3 (14)	WH2-DAD-CT	$K_D \sim 2-3 \mu M$ 50 mM KCl	yes/yes	INF2 C-term interferes mDia1 C-term does not interfere	inhibits elongation (~ nM)
<i>Mm</i> Dia1 (13)	DAD-CT	$K_D \sim 100 \mu M$ 200 mM NaCl	yes/yes	profilin does not interfere	accelerates elongation (~ μM)
<i>Dm</i> Capu (15)	tail	$K_D \sim 20 \mu M$ 50 mM NaCl	no/yes	WH2, RPEL1 interfere profilin does not significantly interfere	inhibits elongation (> 10 μM)
<i>Dm</i> DAAM (in this study)	DAD-CT	$K_D = 44.4 \pm 2.85 \mu M$ 50 mM NaCl	no/yes	profilin does not interfere WH2 interferes	inhibits elongation (> 40-50 μM)

¹The main actin interacting element of the C-terminal regions are highlighted in bold.

FIGURE LEGENDS

Figure 1. Domain organization and sequence alignments of DAAM.

(A) Domain organization of full length *Drosophila melanogaster* DAAM formin (*Dm* DAAM) and of the constructs investigated in this study. G: GTPase binding domain, DID: diaphanous inhibitory domain, DD: dimerization domain, CC: coiled coil, FH: formin homology domain, D: diaphanous autoregulatory domain, CT: C-terminal sequence element. Numbers indicate the number of the first and the last residue in each construct. The positions of the mutated amino acids are highlighted by asterisks. The figure was made with Illustrator for Biological Sciences (57). (B) Sequence alignment of FH2 domains from different formins. The conserved Ile in the FH2 domain is highlighted in red, its position in each formin is given in parenthesis at the end of the sequences. The α D helix of the knob region is underlined. (C) Alignment of C-terminal regions from different formins and WH2 domain sequences. The hydrophobic amino acid triplet and the LKKT/V motif are shown by bold and bold italics, respectively. Positively charged Arg and Lys residues in the CT region are shown in italics. The Arg residues replaced by Ala in the DAAM DAD-CT^{R-A} construct are highlighted in red. Residues that were shown to be important for actin interaction in mDia1, FMNL3 and Capuccino are underlined (13,14,47). UniProt accession numbers are as follows: *Dm* DAAM Q8IRY0, *Hs* DAAM1 Q9Y4D1, *Hs* DAAM2 Q86T65, *Hs* Diaph1 O60610, *Hs* Diaph2 O60879, *Hs* Diaph3 Q9NSV4, *Hs* Fmn1 Q68DA7, *Hs* Fmn2 Q9NZ56, *Hs* FMNL1 O95466, *Dm* Capu Q24120, *Sc* Bnr1 P40450, *Sc* Bni1 P41832, *Hs* WASP P42768, *Hs* WAVE2 Q9Y6W5, *Hs* Lmod1 P29536, *Hs* Lmod2 Q6P5Q4, *Hs* Lmod3 Q0VAK6, *Mm* FMNL3 Q6ZPF4, *Hs* INF2 Q27J81. *Dm*: *Drosophila melanogaster*, *Hs*: *Homo sapiens*, *Sc*: *Saccharomyces cerevisiae*, *Mm*: *Mus musculus*.

Figure 2. cDAAM is more efficient in catalyzing actin assembly than FH1-FH2.

(A) Representative pyrenyl traces of spontaneous and FH1-FH2 or cDAAM catalyzed assembly of free G-actin and profilin:G-actin, as indicated. Final concentrations: [actin] = 2 μ M, [profilin] = 6 μ M, [FH1-FH2] = 200 nM, [cDAAM] = 200 nM. (B) FH1-FH2 and cDAAM concentration dependence of the relative polymerization rate of free G-actin and profilin:G-actin, as indicated. Error bars: standard deviations, $n = 3 - 5$. (C) Steady-state anisotropy of Alexa488NHS-G-actin in the absence and presence of profilin as the function of DAAM concentration, as indicated. Dashed lines in the corresponding colors show the fits to the data according to Eq. 3. Dissociation equilibrium constants are summarized in Table 1. Error bars: standard deviations, $n = 2 - 3$. Final concentrations: [actin] = 0.2 μ M, [LatA] = 4 μ M, [profilin] = 0.8 μ M, [NaCl] = 5 mM. (D) TIRFM montages of actin assembly and representative skeletonized images showing the field of view of actin assembly in the absence and presence of FH1-FH2 or cDAAM, as indicated. Scale bar = 10 μ m, time = min : s. Final concentrations: [actin] = 0.5 μ M, [profilin] = 2 μ M, [FH1-FH2] = 100 nM, [cDAAM] = 100 nM. (E) The number of actin filaments nucleated spontaneously or in the presence of FH1-FH2 and cDAAM derived from skeletonized images. Final concentrations as in panel (D), $n = 20 - 62$. (F) The elongation rate of individual actin filaments polymerized from free, and profilin:G-actin in the absence and presence of FH1-FH2 or cDAAM. Final concentrations as in panel (D), $n = 30 - 89$.

Figure 3. Interactions of DAD-CT and DAD with actin.

(A) Steady-state anisotropy of Alexa488NHS labelled G-actin in the absence and presence of profilin as the function of DAD-CT, DAD and DAD-CT^{R-A} concentration, as indicated. Dashed lines in the corresponding colors show the fits to the data according to Eq. 3. Dissociation equilibrium constants are summarized in Table 1. Error bars: standard deviations, $n = 2 - 4$. Final concentrations: [actin] = 0.2 μ M, [LatA] = 4 μ M, [profilin] = 0.8 μ M, [NaCl] = 5 mM (circles), 17 mM (squares), 50 mM (diamonds). FL: fluorescently labelled protein. (B) Steady-state anisotropy of Alexa568NHS labelled profilin in the presence of G-actin as the function of DAD-CT concentration, as indicated. Dashed line in the corresponding color shows the fit to the data according to Eq. 3. Dissociation equilibrium constant is given in Table 1. Error bars: standard deviations, $n = 2 - 3$. Final concentrations: [actin] = 4 μ M, [LatA] = 8 μ M, [profilin] = 2 μ M, [NaCl] = 5 mM. FL: fluorescently labelled protein. (C) Steady-state anisotropy of Alexa488NHS labelled G-actin in complex with DAD-CT as a function of SALS-WH2 concentration. As controls, the steady-state anisotropy of G-actin in the absence of any binding proteins

and G-actin saturated with SALS-WH2 (1.5 μM) are shown. Final concentrations: [actin] = 0.2 μM , [LatA] = 4 μM , [DAD-CT] = 20 μM , [NaCl] = 5 mM. FL: fluorescently labelled protein. (D) Representative polymerization kinetics of free G-actin and profilin:G-actin in the absence and presence of the C-terminal regions of DAAM, as indicated. Final concentrations: [actin] = 2 μM , [profilin] = 6 μM . (E) TIRFM montages of actin assembly and representative skeletonized images showing the field of view of actin assembly in the absence and presence of DAD-CT, as indicated (for spontaneous actin assembly see *Figure 2, D*). Scale bar = 10 μm , time = min : s. Final concentrations: [actin] = 0.5 μM , [profilin] = 2 μM , [DAAM] = 42 μM . (F) The number of actin filaments nucleated spontaneously or in the presence of DAD-CT derived from skeletonized images. Final concentrations as in panel (E), $n = 16 - 20$. (G) The elongation rate of individual actin filaments polymerized from free, and profilin:G-actin in the absence and presence of DAD-CT (42 μM), DAD (42 μM) or DAD-CT^{R-A} (42 μM). Final concentrations as in panel (E), $n = 24 - 79$.

Figure 4. The effects of DAAM DAD and CT regions in the presence of the native FH2 domain.

A) Representative pyrenyl traces of spontaneous and cDAAM^{R-A} or cDAAM Δ CT catalyzed assembly of free G-actin and profilin:G-actin, as indicated. The data for FH1-FH2 and cDAAM from *Figure 2, E* are shown here as controls. Final concentrations: [actin] = 2 μM , [profilin] = 6 μM , [cDAAM^{R-A}] = 200 nM, [cDAAM Δ CT] = 200 nM. (B) cDAAM^{R-A} or cDAAM Δ CT concentration dependence of the relative polymerization rate of free G-actin and profilin:G-actin, as indicated. Error bars: standard deviations, $n = 3 - 4$. Data obtained for FH1-FH2 and cDAAM from *Figure 2, B* are shown. (C) TIRFM montages of actin assembly and representative skeletonized images showing the field of view of actin assembly in the absence and presence of cDAAM^{R-A} or cDAAM Δ CT, as indicated (for spontaneous actin assembly see *Figure 2, D*). Scale bar = 10 μm , time = min : s. Final concentrations: [actin] = 0.5 μM , [profilin] = 2 μM , [cDAAM^{R-A}] = 100 nM, [cDAAM Δ CT] = 100 nM. (D) The number of actin filaments nucleated spontaneously or in the presence of cDAAM^{R-A} or cDAAM Δ CT derived from skeletonized images. Final concentrations as in panel (C), $n = 20 - 54$. (E) The elongation rate of individual actin filaments polymerized from free, and profilin:G-actin in the absence and presence of cDAAM^{R-A} or cDAAM Δ CT. Final concentrations as in panel (C), $n = 39 - 79$.

Figure 5. The effects of DAAM DAD and CT regions on the loss-of-function mutation of FH2.

(A) Steady-state anisotropy of Alexa488NHS labelled G-actin in the absence and presence of profilin as the function of DAAM concentration, as indicated. The data for FH1-FH2 and cDAAM from *Figure 2, C* are shown here as controls. Dashed lines in the corresponding colors show the fits to the data according to *Eq. 3*. Dissociation equilibrium constants are summarized in *Table 1*. Error bars: standard deviations, $n = 2 - 3$. Final concentrations: [actin] = 0.2 μM , [LatA] = 4 μM , [profilin] = 0.8 μM , [NaCl] = 5 mM. (B) Representative polymerization kinetics of free G-actin and profilin:G-actin in the absence and presence of different regions of DAAM, as indicated. Final concentrations: [actin] = 2 μM , [profilin] = 6 μM . (C) TIRFM montages of actin assembly and representative skeletonized images showing the field of view of actin assembly in the absence and presence of FH1-FH2^{I732A} and cDAAM^{I732A}, as indicated (for spontaneous actin assembly see *Figure 2, D*). Scale bar = 10 μm , time = min : s. Final concentrations: [actin] = 0.5 μM , [profilin] = 2 μM , [FH1-FH2^{I732A}] = 100 nM, [cDAAM^{I732A}] = 100 nM. (D) The number of actin filaments nucleated spontaneously or in the presence of FH1-FH2^{I732A} and cDAAM^{I732A} derived from skeletonized images. Final concentrations as in panel (C), $n = 20 - 50$. (E) The elongation rate of individual actin filaments polymerized from free, and profilin:G-actin in the absence and presence of FH1-FH2^{I732A} and cDAAM^{I732A}. Final concentrations as in panel (C). 20 μM and 2 μM indicate the data obtained in the presence of higher concentrations of FH1-FH2^{I732A} and cDAAM^{I732A}, respectively, $n = 15 - 79$. (F) Critical concentration (J(c)) plots of actin assembly in the absence and presence of cDAAM^{I732A} (1 μM), as indicated. Inset: enlarged view of the initial part of the J(c) plot, red arrow highlights the breaking points corresponding to the unassembled actin at steady-state. Dashed lines in the corresponding colors show the fit to the linear part of the data. Error bars: standard deviations, $n = 3$.

Figure 6. The antagonism between DAAM and CP.

(A) Polymerization of profilin:G-actin initiated in the absence and presence of DAAM constructs, as indicated. The addition of CP after 120 s is indicated by arrow. Final concentrations: [actin] = 2 μ M, [profilin] = 6 μ M, [CP] = 68 nM, [DAAM] = 200 nM. (B) Polymerization rates before ([CP] = 0 nM) and after addition of CP at different concentrations derived from pyrenyl traces. Dashed lines in the corresponding color show the fit of the data using Eq. 2. The fit gave IC₅₀ values of IC₅₀(cDAAM) = 345.9 \pm 27.60 nM, IC₅₀(FH1-FH2) = 47.7 \pm 16.97 nM, IC₅₀(cDAAM Δ CT) = 108.6 \pm 19.35 nM and IC₅₀(cDAAM^{R-A}) = 93.7 \pm 19.06 nM for CP inhibition. Error bars: standard deviations, n = 2 – 4.

Figure 7. The interaction of DAD-CT with actin filaments.

(A) The DAAM : F-actin ratio in the pellet (P) derived from the analysis of SDS-PAGE gels. Dashed lines correspond to the fit of the data (21,35), the fit gave K_D values of K_d(DAD-CT) = 38.9 \pm 3.2 μ M, K_D(DAD) > 100 μ M. [F-actin] = 2.5 μ M, Error bars: standard deviations, n = 2 – 4. (B) Bundling activity of different regions of DAAM. The relative amount of actin filaments in the supernatant (SN) as the function of DAAM concentration, determined from the quantification of SDS-PAGE analysis of the samples. Final concentrations: [F-actin] = 1 μ M. Error bars: standard deviations, n = 2 – 3.

Figure 8. The FH2:actin interaction is regulated through the DID:DAD-CT interaction.

(A) Representative polymerization kinetics of actin in the absence and presence of DID, FH1-FH2 and cDAAM, as indicated. [actin] = 2 μ M (containing 5 % pyrenyl actin), [FH1-FH2] = 5 μ M, [cDAAM] = 0.51 μ M. (B) The relative polymerization rate of cDAAM catalyzed actin assembly as a function of DID concentration. Dashed line corresponds to the fit of the data using Eq. 1.. The fit gave half-inhibition of cDAAM by DID at 30.5 \pm 14.31 nM. The relative rate of spontaneous actin assembly is indicated by grey dotted line.

Figure 9. Constitutively active dDAAM induces filopodia formation in *Drosophila* primary neurons.

(A) Domain organization of full length *Drosophila melanogaster* DAAM formin and of the constructs investigated in the *in vivo* experiments. Abbreviations as in Figure 1A. The position of the I732A mutation is highlighted by asterisk. The figure was made with Illustrator for Biological Sciences (57). (B) Representative images of primary neurons obtained from control or constitutively active dDAAM overexpressing embryos, stained with tubulin (magenta), actin (green) and HRP (red). Scale bar = 5 μ m. (C) Expression of Δ DAD-dDAAM (*c155>UAS- Δ DAD-dDAAM*, n = 62) construct induced a significant increase of filopodia number compared to control (*c155/+*, n = 162) and Δ DAD-CT-dDAAM I732A (*c155>UAS- Δ DAD-dDAAM I732A*, n = 60) expressing cells. The other genotypes were not significantly different (*c155> Δ DADCT-dDAAM*, n = 61; *c155> Δ DADCT-dDAAM I732A*, n = 68). Data presented on the figure are given as mean \pm SEM. Data were analyzed with Kruskal-Wallis for the whole data set, followed by Dunn's *posthoc* test for multiple comparison. (D) Representative image from a time lapse sequence of a neuronal growth cone expressing Actin5C::GFP. Scale bar = 5 μ m (corresponding to Supplemental movie 1). (E) Comparison of filopodia extension rate showed no significant difference (One way analysis of variance, p = 0.09; Control: 5.4 \pm 1.2, mean \pm SD, n=3/134; Δ DAD-dDAAM: 3.6 \pm 1.2, mean \pm SD, n=4/118; Δ DAD-CT-dDAAM: 4.8 \pm 0.3, mean \pm SD, n=4/77). (F) The retraction rate is significantly lower in the case of Δ DAD and Δ DAD-CT overexpression, compared to the control filopodia (One way analysis of variance, p = 0.0047; Control: 6.4 \pm 0.7, mean \pm SD, n=3/91; Δ DAD-dDAAM: 3.6 \pm 0.5, mean \pm SD, n=4/63; Δ DAD-CT-dDAAM: 4.5 \pm 0.9, mean \pm SD, n=4/40).


Figure 10. Structural model of the interactions of FH2, DAD and CT with actin.

(A) The model was generated using X-ray structures of complexes of FH2:actin and WH2:actin. Four actin subunits (gray, indicated by numbers) are shown according to their arrangement in the Oda's model (58). The FH2 dimer of DAAM (blue) and the DAD region of mDia1 (red) are represented as ribbons. Red dashed lines indicate the possible orientation of the disordered CTs of DAAM. Blue dashed lines indicate the ~ 20 aa linkers connecting the FH2 and DAD of DAAM. Distances are given in Å. The model was generated with PyMOL based on the alignment of following structures: PDB IDs: 4EAH ((48) FMNLFH2-TMR-actin), 2Z6E ((59) hDAAM-FH2), 2BAP ((60) mDia1-DAD), 2A41 ((61) WIP-WH2). (B) Alternative model describing actin nucleation mediated by FH2-DAD-CT for low-affinity

The C-terminus of formins in actin dynamics

C-terminal formin regions. In this scenario, the binding of DAD-CT to actin requires increased affinity, which may be manifested through monomer capturing by FH2.

Figure 1



B FH2 domains

			::	*	:	:	:		
<i>Dm</i>	DAAM	701	DGSYEDLRVTGKAAKQKVL	SVIDGRR	AONCT	I	LLSKLK	-MSDMEISKAILSMD	SNEQLQLD	760	(I732)
<i>Hs</i>	DAAM1	669	ADAIDDTLSS--KLKVKEL	SVIDGRR	AONCN	I	LLSRLK	-LSNDEIKRAILTM	DEQEDLPKD	726	(I698)
<i>Hs</i>	DAAM2	654	LGSTEDIYLA--SRKVKEL	SVIDGRR	AONCI	I	LLSKLK	-LSNEEIRQAILKM	DEQEDLAKD	711	(I683)
<i>Hs</i>	Diaph1	831	DQEGGEEKSVQKKVKEL	KVLDSKTA	QNLS	I	FLGSFR	-MPYQEIKNVILEV	NEA-VLTES	889	(I845)
<i>Hs</i>	Diaph2	690	ALE--EKKTGPTKKVKEL	RILDPKTA	QNLS	I	FLGSYR	-MPYEDIRNVILEV	NED-MLSEA	746	(I720)
<i>Hs</i>	Diaph3	698	DIE--EKKS--IKKIKEL	KFLDSKIA	QNLS	I	FLSSFR	-VPYEEIRMMILEV	DET-RLAES	752	(I705)
<i>Hs</i>	Fmn1	1037	TYEK-----KNKVKKI	IKLLDGKRS	QTVG	I	LISSLH	-LEMKDIQQAIFNV	DDS-VVDLE	1088	(I1061)
<i>Hs</i>	Fmn2	1347	TIS-----KTKAKQV	VKLLSNKRS	QAVG	I	LMSSLH	-LDMKDIQHAVVN	LDNS-VVDLE	1397	(I1208)
<i>Hs</i>	FMNL1	689	SLDLSALKSKAAQKAP	SKATLIEAN	RAKNLA	I	TLRKGN	-LGAERICQAIEAY	DLQ-ALGLD	747	(I720)
<i>Dm</i>	Capu	681	PKEL-----KVKRAK	SIKVLDPERS	RNVG	I	IWRSLH	-VPSSEIEHAIYH	IDTS-VVSLE	733	(I706)
<i>Sc</i>	Bnr1	936	SSIALSSNNGKSSNEL	KKISFLSRD	LAQQFG	I	NLHMFS	QLSDMEFVMKVLN	CNDNDIVQNVN	996	(I967)
<i>Sc</i>	Bni1	1412	-----KEDLQKITF	LSRDISQ	QFG	I	NLHMYSSL	SVADLVKKILNC	DRDFLQTPS	1460	(I1431)

C DAD | CT regions – WH2 domains

			* : ::					
Dm	DAAM	1084	GDNKGEFDD	L I S A L R T G	DVFGEDMAKFKRS-		R K A R V L N G G G S S T G H T S P P R H G S L Q R E E S G R E R E R T V R R Q -----	1153
Hs	WASP	430	-----	G R G A L L D Q I R Q G --	I Q L N K T - -----		-----	447
Hs	WAVE2	436	-----	A R S D L L S A I R Q G --	F Q L R R V - -----		-----	453
Hs	Lmod1	574	-----	S R D Q L L A A I R S S N L K Q L K K V - -----	-----		-----	593
Hs	Lmod2	521	-----	A H E N L M E A I R G S S I K Q L K R V - -----	-----		-----	540
Hs	Lmod3	534	-----	P R D Q L L N D I R H S S V A Y L K P V - -----	-----		-----	553
Mm	FMNL3	967	-----	Q Q E L I A E L L R R R - --	Q A K E H - -----		-----	982
Hs	INF2	974	-----	D A L L A D I R K G --	F Q L R K T - -----		-----	989
Mm	Diaph1	1194	-DETGV	MDS L L E A L Q S G A A F -	R R K R G P R Q A N		RKAGCAVTSLLASELTKDDAMAAVPAKVSKNSETFPTILEEAKELVGRAS	1272
Dm	Capu	1033					--- R R N -----V S T K V E K S G R I S L K E R M L M R R S K N -----	1059

Figure 2




Figure 3



Figure 4




Figure 5




Figure 6




Figure 7







Figure 8

A



B






Figure 9


A




C




D



B



E



F






Figure 10

A



B



The activities of the c-terminal regions of the formin protein disheveled-associated activator of morphogenesis (daam) in actin dynamics

Andrea Teréz Vig, István Földi, Szilárd Szikora, Ede Migh, Rita Gombos, Mónika Ágnes Tóth, Tamás Huber, Réka Pintér, Gábor Csaba Talián, József Mihály and Beáta Bugyi

J. Biol. Chem. published online June 22, 2017

Access the most updated version of this article at doi: [10.1074/jbc.M117.799247](https://doi.org/10.1074/jbc.M117.799247)

Alerts:

- [When this article is cited](#)
- [When a correction for this article is posted](#)

[Click here](#) to choose from all of JBC's e-mail alerts

Supplemental material:

<http://www.jbc.org/content/suppl/2017/06/22/M117.799247.DC1>

This article cites 0 references, 0 of which can be accessed free at

<http://www.jbc.org/content/early/2017/06/22/jbc.M117.799247.full.html#ref-list-1>